

U. S. DEPARTMENT OF AGRICULTURE

BUREAU OF SOILS

IN COOPERATION WITH  
WISCONSIN GEOLOGICAL AND NATURAL HISTORY SURVEY  
AND UNIVERSITY OF WISCONSIN COLLEGE OF AGRICULTURE

---

# SOIL SURVEY OF GREEN LAKE COUNTY WISCONSIN

BY

W. J. GEIB, IN CHARGE, A. C. ANDERSON, E. H. BAILEY, AND  
M. J. EDWARDS, U. S. DEPARTMENT OF AGRICULTURE, AND  
HOMER CHAPMAN, OSCAR MAGISTAD, F. J.  
O'CONNELL, T. J. DUNNEWALD, AND  
KENNETH WHITSON, WISCONSIN  
GEOLOGICAL AND NATURAL  
HISTORY SURVEY

---

[Advance Sheets—Field Operations of the Bureau of Soils, 1922]



UNITED STATES  
GOVERNMENT PRINTING OFFICE  
WASHINGTON  
1928

BUREAU OF SOILS

MILTON WHITNEY, *Chief*  
WILBERT W. WEIR, *Acting Editor*

SOIL SURVEY

CURTIS F. MARBUT, *Chief*  
MARK BALDWIN, *Inspector, District No. 1*

---

COOPERATION

WISCONSIN GEOLOGICAL AND NATURAL HISTORY SURVEY

ERNEST F. BEAN, *Director*

AND

UNIVERSITY OF WISCONSIN, COLLEGE OF AGRICULTURE

H. L. RUSSELL, *Dean*

A. R. WHITSON, *In Charge Soil Survey*

U. S. DEPARTMENT OF AGRICULTURE

BUREAU OF SOILS

IN COOPERATION WITH  
WISCONSIN GEOLOGICAL AND NATURAL HISTORY SURVEY  
AND UNIVERSITY OF WISCONSIN COLLEGE OF AGRICULTURE

---

# SOIL SURVEY OF GREEN LAKE COUNTY WISCONSIN

BY

W. J. GEIB, IN CHARGE, A. C. ANDERSON, E. H. BAILEY, AND  
M. J. EDWARDS, U. S. DEPARTMENT OF AGRICULTURE, AND  
HOMER CHAPMAN, OSCAR MAGISTAD, F. J.  
O'CONNELL, T. J. DUNNEWALD, AND  
KENNETH WHITSON, WISCONSIN  
GEOLOGICAL AND NATURAL  
HISTORY SURVEY

---

[Advance Sheets—Field Operations of the Bureau of Soils, 1922]



UNITED STATES  
GOVERNMENT PRINTING OFFICE  
WASHINGTON  
1928

[PUBLIC RESOLUTION—No. 9.]

JOINT RESOLUTION Amending public resolution numbered eight, Fifty-sixth Congress, second session, approved February twenty-third, nineteen hundred and one, "providing for the printing annually of the report on field operations of the Division of Soils, Department of Agriculture."

*Resolved by the Senate and House of Representatives of the United States of America in Congress assembled*, That public resolution numbered eight, Fifty-sixth Congress, second session, approved February twenty-third, nineteen hundred and one, be amended by striking out all after the resolving clause and inserting in lieu thereof the following:

That there shall be printed ten thousand five hundred copies of the report on field operations of the Division of Soils, Department of Agriculture, of which one thousand five hundred copies shall be for the use of the Senate, three thousand copies for the use of the House of Representatives, and six thousand copies for the use of the Department of Agriculture: *Provided*, That in addition to the number of copies above provided for there shall be printed, as soon as the manuscript can be prepared, with the necessary maps and illustrations to accompany it, a report on each area surveyed, in the form of advance sheets, bound in paper covers, of which five hundred copies shall be for the use of each Senator from the State, two thousand copies for the use of each Representative for the congressional district or districts in which the survey is made, and one thousand copies for the use of the Department of Agriculture.

Approved, March 14, 1904.

[On July 1, 1901, the Division of Soils was reorganized as the Bureau of Soils.]



# CONTENTS

	Page
County surveyed.....	1801
Climate .....	1804
Agriculture.....	1805
Soils .....	1807
Parr silt loam.....	1810
Parr loam.....	1812
Parr fine sandy loam.....	1813
Dodgeville silt loam.....	1814
Bellefontaine fine sandy loam, dark-colored phase.....	1814
Bellefontaine sandy loam, dark-colored phase.....	1815
Bellefontaine loam.....	1817
Bellefontaine silt loam.....	1817
Bellefontaine gravelly loam.....	1818
Bellefontaine gravelly sandy loam.....	1819
Miami silt loam.....	1819
Miami loam.....	1821
Coloma sand.....	1822
Coloma fine sand.....	1822
Warsaw loam.....	1823
Warsaw fine sandy loam.....	1823
Warsaw silt loam.....	1824
Fox sandy loam.....	1824
Fox loam.....	1825
Fox silt loam.....	1825
Plainfield sand.....	1826
Plainfield fine sand.....	1827
Superior fine sandy loam.....	1828
Maumee silt loam.....	1828
Maumee loam.....	1829
Clyde silt loam.....	1830
Clyde loam.....	1830
Poygan silt loam.....	1831
Poygan clay loam.....	1831
Dunning fine sand.....	1831
Genesee silt loam.....	1832
Peat .....	1832
Rough stony land.....	1836
Suggestions for soil improvement.....	1836
Lime and fertilizers.....	1836
Crop rotations .....	1838
Summary.....	1840

# ILLUSTRATIONS

## PLATE

	Page
PLATE L, Fig. 1.—Hemp mill at Markesan, Wis. Fig. 2.—An area of Clyde silt loam used for pasture.....facing	1828

## FIGURE

Fig. 54.—Sketch map showing location of Green Lake County, Wis.....	1801
---	------

## MAP

Soil map, Green Lake County, Wis.



# SOIL SURVEY OF GREEN LAKE COUNTY, WIS.

By W. J. GEIB, in Charge, A. C. ANDERSON, E. H. BAILEY, and M. J. EDWARDS, United States Department of Agriculture, and HOMER CHAPMAN, OSCAR MAGISTAD, F. J. O'CONNELL, T. J. DUNNEWALD, and KENNETH WHITSON, Wisconsin Geological and Natural History Survey

## COUNTY SURVEYED

Green Lake County lies slightly southeast of the center of Wisconsin, and the city of Green Lake, the county seat, is 90 miles by rail from Milwaukee. The greatest length north and south is 24 miles, the greatest width east and west is 18 miles, and its area is 360 square miles, or 230,400 acres.

One of the most interesting surface features of Green Lake County and economically considered, by far the most important, is the extensive area of prairie land which occurs as a high plain or plateau in the southeastern part, at an average elevation of about 1,000 feet above sea level. The surface of this plain is for the most part level or gently rolling. It is confined chiefly to Green Lake, Mackford, and Manchester Townships, and includes some of the best farming land in the county.

This rolling plain extends northward to an imaginary line drawn northeast and southwest through Center House, where there is a sharp drop into the valley occupied by Green Lake. This lake has a maximum depth of 235 feet and the surface water level is 796 feet above sea level, which makes the lake bottom or valley floor, between 400 and 500 feet lower than the high plain. This valley or trough which has a sandstone floor, extends in a northeast-southwest direction and at one time probably joined Puckaway Lake, but it is now separated from the marshes bordering that lake by a marginal moraine about one-half mile wide.

North of Green Lake is another elevation similar to the one south of the lake, but nearly 100 feet lower. The bedrock north of the lake is chiefly magnesian limestone, and that south of the lake is Trenton limestone. The drop from the limestone formations forms an escarpment with steep slopes which is a conspicuous feature of the landscape. North of Green Lake this drift-covered magnesian escarpment extends westward almost to Princeton and then swings northeastward to Berlin, and north and east of this general line the land is rolling. Mount Tom in section 11, T. 16 N., R. 12 E., is a



FIG. 54.—Sketch map showing location of Green Lake County, Wis.

prominent hill of sandstone capped with limestone, with an elevation of more than 150 feet above the adjoining lowland.

The northwestern part of the county forms part of the central plain of Wisconsin. It is a low, nearly level region of sandy soils and marshland drained by Fox River and its tributaries, and has an elevation ranging from 760 to about 800 feet. South of the extensive marshes which border Lake Puckaway the land is gently rolling and the soils are loamy.

The county lies almost entirely within the Fox River drainage basin. The only exception to this is part of one township in the extreme southeastern corner of the county which drains into Rock River. The main channel of Fox River passes through Lake Puckaway in the southwestern part of the county, flows north and north-east, and leaves the county near Berlin. Beyond the county line it flows into Lake Winnebago, and thence into Green Bay.

Grand River drains much of the southern part of the county including more than one-half the prairie region, and has cut a deep ravine through Mackford, Manchester, and Kingston Townships. It flows westward and joins Fox River in Marquette County not far from Puckaway Lake. The east-central part of the county drains into Green Lake, thence through Puchyan River into Fox River. The tributary streams have considerable fall, but the current in Fox River is sluggish. In upper Fox River, which includes that part of the river flowing into Lake Winnebago, the fall is less than 6 inches to the mile.

Fox River is interrupted in its course by Lake Buffalo and Lake Puckaway. Lake Buffalo is a shallow crescent-shaped lake in Marquette County, about 12 miles long and three-fourths mile wide; and Lake Puckaway, partly in Green Lake County and partly in Marquette County, is 7 miles long and 1½ miles wide. The rock floor of the preglacial valley at Lake Puckaway is buried to a depth of more than 300 feet. Extensive swamps border Fox River, and these swamps and lakes regulate the flow of lower Fox River and prevent or diminish floods.

The elevation of certain points in Green Lake County, as compiled from the records of the Geological Survey, and from other sources is given below:

*Elevation of different points in Green Lake County*

Place	Authority	Elevation
		<i>Feet</i>
Markesan <sup>1</sup> .....	Chicago, Milwaukee & St. Paul Railway.....	859
Green Lake, surface of water.....	United States Geological Survey.....	796
Green Lake (city) <sup>1</sup> .....	Chicago & North Western Railway.....	902
Princeton <sup>1</sup> .....	do.....	881
Berlin <sup>1</sup> .....	Chicago, Milwaukee & St. Paul Railway.....	768
Lake Puckaway.....	United States Geological Survey.....	764
Saint Marie <sup>1</sup> .....	Chicago & North Western Railway.....	895
School No. 8, Green Lake Township.....	United States Geological Survey.....	1,024
Searles School, Green Lake Township.....	do.....	1,014
Dalton.....	Chicago & North Western Railway.....	824

<sup>1</sup> GANNETT, H., A DICTIONARY OF ALTITUDES IN THE UNITED STATES. U. S. Geol. Survey Bul. 274, 1072 p. 1906. (Ed. 4.)

The following table shows the population of the county and of several of the communities within the county as reported by the census for the years 1900, 1910, and 1920:

*Population of Green Lake County for three census years*

Place	1920	1910	1900
Green Lake County.....	14, 875	15, 491	15, 797
Berlin (city).....	4, 356	4, 586	4, 448
Green Lake (village).....	456	563	450
Princeton (village).....	1, 275	1, 269	1, 202
Markesan (village).....	959	892	706

In the census of 1920 the city of Berlin was the only community classed as urban and the remainder of the population, or 70.7 per cent, was classed as rural. The density of the rural population was 29.2 persons to the square mile. The distribution is fairly uniform, although the most thickly settled regions are on the better soils in the southern, eastern, and northeastern parts of the county.

Green Lake County was formed from territory which was previously a part of Marquette County and was organized in 1858, and the present boundaries were defined in 1860. Green Lake County was named for its principal body of water, the lake being named by the French "Lac Verd," which the early settlers translated into Green Lake. Among the early settlers were many Germans and English, and some Irish, Poles, and Scandinavians. Many early settlers came also from the older Eastern States. The first settlement was made in this region some time between 1830 and 1835.

The county is served by two railway systems which provide fairly good facilities for freight shipment and passenger traffic. The new Short Line of the Chicago & North Western Railway between Chicago and the Twin Cities, crosses the extreme southwestern corner of the county and passes through Dalton. A branch of this system extending from Milwaukee through Fond du Lac to Marshfield crosses the county from east to west and passes through Green Lake and Princeton. A branch of the Chicago, Milwaukee & St. Paul Railway enters the southeastern part of the county, extends as far as Markesan, and connects at Brandon in Fond du Lac County with the branch from Ripon to Milwaukee. Another line of the same system connects Ripon in Fond du Lac County with Berlin in the northeastern part of Green Lake County.

The country roads are undergoing rapid improvements, and several State and county trunk highways also traverse the county. The materials used in improving the roads consist of concrete, crushed rock, and gravel. A concrete road connects Green Lake and Berlin and another one extends part way from Green Lake to Princeton, and many roads are macadamized. The dirt roads in the northern and western parts of the county are sandy in places, but throughout these sections, as elsewhere, main trunk highways are under construction. All parts of the county are reached by rural mail routes and telephone communication is available everywhere.

The towns do not afford a market for all the farm produce, but they supply shipping facilities which render the markets of Milwau-



kee and Chicago very accessible. Much of the marketing is done through cooperative shipping associations.

#### CLIMATE<sup>1</sup>

Practically all of Green Lake County lies within the Fox River drainage basin and forms part of the Fox and Wolf River basin which is one of the eight climatic provinces in Wisconsin. The climate of this basin is influenced somewhat by lakes, but it possesses more of the characteristics of a land climate. The winters at Green Bay, Appleton, Pine River, Oshkosh, and Fond du Lac, with a mean temperature averaging 18.7° F., are as warm as those at Dodgeville or Lancaster; and spring and summer are as cool as the average at Eau Claire and Osceola. The frost-free season which varies from 130 to 150 days, shows land influence, lasting about the same length of time as the frost-free season in Richland, Buffalo, and St. Croix Counties, Wis.; Cattaraugus, Chenango, and Delaware Counties, N. Y.; Center and Lycoming Counties Pa.; northern Iowa; or central Utah. The rainfall in this region is possibly a little less than elsewhere in the State.

The mean annual temperature for Green Lake County is 44.9° F. as recorded at the nearest Weather Bureau station at Grand River Locks in Marquette County. The maximum recorded temperature is 103° and the minimum -40°. The average rainfall is 33.26 inches, which under normal conditions is well distributed throughout the growing season. April, May, June, July, August, and September, during which the crops make most of their growth and are most in need of a good supply of moisture, average 3.86 inches of rainfall. Although the range in temperature is wide, the extremely low and extremely high readings recorded are rare and the periods are of short duration. The winters are severe with an average snowfall of 43.1 inches. The winds are usually from the northeast.

The average date of the last killing frost in spring is May 5, and of the first in the fall, September 30, giving an average frost-free season of 147 days. The latest recorded frost in the spring occurred on May 23, and the earliest recorded frost in the fall on September 10.

The climatic conditions are representative of a large section in the east-central and central parts of the State, where agriculture is highly developed on all of the best soils and where extremes of climate do not prevent the growing of any general farm crops. Excellent water for household purposes and for stock is available throughout the county. This fact, together with healthful climatic conditions, a long growing season, and large areas of rich, productive soil, helps to make Green Lake County one of the most desirable agricultural communities of Wisconsin.

The following table compiled from the records of the Weather Bureau station at Grand River Locks, Marquette County, gives the more important climatic data representative of this county:

<sup>1</sup> For more information on climate and its relation to agriculture see the following bulletin, on which this chapter is based: WHITSON, A. R., and BAKER, O. E. THE CLIMATE OF WISCONSIN AND ITS RELATION TO AGRICULTURE. Wis. Agr. Expt. Sta. Bul. 223, 65 p., illus. 1912.

*Normal monthly, seasonal, and annual temperature and precipitation at Grand River Locks, Marquette County*

(Elevation, 768 feet)

Month	Temperature			Precipitation			
	Mean	Absolute maximum	Absolute minimum	Mean	Total amount for the driest year (1923)	Total amount for the wettest year (1906)	Snow, average depth
	° F.	° F.	° F.	Inches	Inches	Inches	Inches
December .....	21.4	57	-24	1.20	1.61	2.22	10.1
January .....	14.5	48	-33	1.15	.79	3.11	11.4
February .....	17.6	53	-40	1.03	.76	1.40	9.0
Winter .....	17.8	57	-40	3.38	3.16	6.73	30.5
March .....	31.9	82	-26	1.87	1.79	2.59	5.2
April .....	45.8	85	14	2.58	2.74	1.35	2.6
May .....	56.8	91	26	4.69	1.27	4.62	.5
Spring .....	44.8	91	-26	9.14	5.80	8.56	8.3
June .....	66.4	99	34	4.33	2.02	9.47	.0
July .....	71.0	103	37	3.76	2.61	2.97	.0
August .....	68.2	98	31	3.77	1.28	6.30	.0
Summer .....	68.5	103	31	11.86	5.91	18.74	.0
September .....	61.0	98	23	4.05	3.78	4.15	.0
October .....	48.8	84	11	2.87	1.73	5.25	.9
November .....	35.7	72	-10	1.96	.80	4.07	3.4
Fall .....	48.5	98	-10	8.88	6.31	13.47	4.3
Year .....	44.9	103	-40	33.26	21.18	47.50	43.1

## AGRICULTURE

The most common form of agriculture in Green Lake County is general farming, with dairying as its most important feature. In 1919 there were 26,688 cattle in the county. Of these, 22,593 were dairy cattle and the remainder were beef cattle. The quantity of milk produced was 4,547,840 gallons. The value of milk, cream, and butterfat actually sold, and butter and cheese made on the farms was \$1,121,001. The average quantity of milk produced was 404 gallons per cow.

A list published by the Wisconsin Livestock Breeders Association showed that in Green Lake County there are 210 breeders of cattle, listed as follows: Holstein breeders, 100; Guernsey breeders, 34; Shorthorn breeders, 40; Brown Swiss breeders, 20; Hereford breeders, 8; Jersey breeders, 4; Red Polled breeders, 4.

Other farmers are raising cattle; in fact, practically every farmer in the county raises some, but comparatively few make a business of raising purebred cattle or high standard grades for breeding purposes. The proportion of purebred livestock is gradually increasing. Raising beef cattle here is not so important an industry as dairying; but many farmers buy steers and fit them for market, and others sell as beef cattle part or all of the cattle they raise.

In 1919 there were 21,887 hogs and 8,223 sheep, the total value being \$357,397 and \$76,282 respectively. The sheep produced 48,035

pounds of wool valued at \$26,318. The value of chickens and eggs produced in 1919 was \$253,080, and the 165,345 chickens produced 468,406 dozen eggs. The income from the sale of eggs and chickens was \$123,597.

The chief crops are hay, oats, corn, rye, wheat, barley, and potatoes; and the minor crops are peas, beans, sugar beets, hemp, flax, buckwheat, vegetables, and small fruits.

In 1919 there were 46,859 acres of hay and forage crops, the average yield being a little more than 2 tons an acre. Of this acreage 17,117 acres were in tame grasses which included 3,702 acres in timothy, 9,939 acres in mixed timothy and clover, 2,320 acres in clover, and 195 acres in alfalfa.

The production of alfalfa is growing in favor, although it is difficult to obtain a good stand because the soil lacks lime and because of winter killing. The acreage in wild hay and marsh grass for hay nearly equals the acreage of tame grasses, but this wild hay is coarse and decidedly inferior to clover and alfalfa. Clover and alfalfa grow best on Miami and Bellefontaine soils, but if lime and fertilizers are used, excellent crops may be raised on all the well-drained soils of the county. Hay is grown on all soils of the county, but does best on the heavier soils.

According to the census there were 27,380 acres in oats in 1919 which yielded 765,311 bushels of grain. This crop is grown in all parts of the county, but the largest acreage is in the southern, southeastern, and eastern parts, where loam and silt loam soils predominate. Corn was grown on 21,415 acres in 1919 yielding 849,208 bushels. It is grown on practically all soils, although the black prairie soils and the well-drained Clyde silt loam give most satisfactory returns. There were 9,920 acres of rye yielding 127,180 bushels, and 7,433 acres of wheat yielding 98,731 bushels which was somewhat less than the acreage during the period of the World War but more than that for some years prior to the war. Wheat is grown chiefly on the Miami and Parr silt loam soils which are well adapted to it. There were 2,052 acres in potatoes which are grown mostly on sandy loams, fine sandy loams, and other fine sandy soils.

Hemp grown for fiber is a comparatively new crop. It is shipped to hemp mills at Markesan, Fair Water, and Ripon. Hemp brought large profits immediately following the World War, but with the price of cotton reduced the price of hemp also declined and the returns were less satisfactory. Hemp requires fertile soil for profitable yields, and at present it is grown mostly on Miami and Parr silt loams. The erection of a mill for handling hemp is an item of considerable expense, but before the growers in any community can raise hemp on a large scale it is necessary that they should have access to a hemp mill. (Pl. L, fig. 1.)

It is recognized that different soils are adapted to different kinds of farming. Gravelly loam soils are in some places too rough for growing general farm crops, and farmers realize that such areas are best suited for pastures. It is well known that crops on lowlands are most susceptible to frost. It is generally conceded that corn grows best on heavy Clyde silt loam, and adapts itself well to all the heavier, dark-colored soils, such as Parr and Warsaw silt loams. On these dark soils which are high in organic matter, small grains are



apt to lodge, and the quality of the grain is not so good as on the light-colored, heavy soils. Peas do best on Miami silt loam and Fox silt loam, and potatoes grow best on sandy and fine sandy loam soils. The sugar content of beets grown on Parr silt loam, Warsaw silt loam, and Clyde silt loam is lower than that of beets grown on Miami silt loam and Fox silt loam, but the yield is sufficiently high to give better returns. The lighter-textured soils are best for truck crops.

Farm buildings and equipment differ greatly throughout the county. In sections in which occur such soils as Parr silt loam and Miami silt loam, which are the heavy light-colored soils, nearly all the farm buildings are large and substantial, and there are many silos. Most of the fences are good, many of them being of woven wire. The work animals are draft horses of medium or heavy weight. Modern farm machinery is used, and tractors for plowing and other farm work are common. In sections where the soil is extremely sandy the buildings, livestock, and machinery are inferior, and where the soils are loamy the farm equipment is of medium grade.

Threshing outfits travel about the county serving the farmers soon after harvest. Several cooperative threshing outfits are owned jointly by groups of farmers, especially where purebred grain is grown to be sold as seed grain.

The tendency throughout Green Lake County is toward better methods of cultivation, fertilization, and seed selection, and as a result, yields are increasing gradually. Where there are sandy soils underlain by sand, spring plowing is preferred to fall plowing. If such soils are covered during fall and early spring with a good growth, loss of plant-food elements by leaching and loss of finer sand particles by severe winds are prevented. Seeding rye in corn rows at the time of last cultivation or in potato fields at digging time prevents much loss of plant-food elements.

When farm work allows the heavy soils, especially those in sod, are plowed in the fall. It is customary to apply stable manure to land which is to be plowed for corn, but if the land has been plowed in the fall the manure is often hauled out during the winter and scattered over the plowed surface. When stubble land is plowed in late summer, manure is frequently applied before plowing. Fall plowing aids in conserving moisture, helps to improve the tilth, and where there is no danger from erosion it is a good practice to follow on heavy soils.

Throughout the county most farmers seed their land to grasses at least once every four or five years. The practice of using lime to correct acidity and to prepare land for clover and alfalfa, is becoming more common. Commercial fertilizers are also being introduced, and, where properly applied, results are satisfactory.

#### SOILS

The soils of Green Lake County are extremely diverse, ranging in texture from heavy clay loam to sand, in composition from nearly pure quartz to highly complex mixtures of many minerals and chemical compounds, and in drainage conditions from very poor to excessive. These soils have general geographic groupings on the map, although they are intermingled in places.

If a line be drawn from the northeastern corner of the county to the southwestern corner, the soils to the west are predominantly sandy, with considerable marshland, and the soils to the east are predominantly heavy, being mostly well-drained silt loam.

The underlying rocks in the eastern part of the county are principally Trenton and Lower Magnesian limestones. The country rock in the western part of the county is Potsdam sandstone. The parent materials of the soils were partly ice laid, partly water laid, and probably partly wind laid. In addition there are soil materials which have been formed through the accumulation of vegetable matter under water, resulting in the formation of peat.

Although there are distinct variations in the soils within Green Lake County, the upland soils, derived from heavy to medium textured drift, are composed of a consistent succession of texture layers, or soil horizons, over extensive areas. These layers or horizons consist of:

(1) A comparatively light-textured surface layer from 6 to 15 inches thick, usually silt loam or loam; (2) a layer of heavier-textured material from 12 to 18 inches thick, commonly called the subsoil; and (3) the partly weathered or unweathered parent material which is lighter in texture than the subsoil. This generalized soil profile, which is the normal profile of the region, is characteristic of the Miami, Bellefontaine, Fox, Parr, and Warsaw soils. An important variation in the color of the surface soils occurs within this group of normally developed soils. The Miami, Bellefontaine, and Fox soils are light grayish brown or brown, and the Parr and Warsaw soils range from dark brown to nearly black. This difference in soil color bears some relationship to the difference in native vegetation, the light-colored soils being forested and the dark-colored soils constituting the prairie lands.

Soils of Green Lake County are classified as 31 types and 6 phases, representing 14 series, not including peat and rough stony land.

Parr soils are dark-colored, well-drained prairie soils derived from calcareous glacial drift. The Parr soils mapped are silt loam with a level phase, loam with a gravelly phase, and fine sandy loam.

The Dodgeville series includes dark-colored, well-drained soils, underlain by limestone. Dodgeville silt loam is the only soil of this series mapped in Green Lake County.

The Bellefontaine soils are similar to those of the Miami series but the surface of the areas is more irregular, the topsoil is browner, the subsoil is brown or red and shows more thorough oxidation, and the parent material is coarser. Bellefontaine soils mapped in Green Lake County are the silt loam, loam, fine sandy loam, dark-colored phase, sandy loam, dark-colored phase, gravelly loam, and gravelly sandy loam.

The Miami series includes light-colored upland soils developed under forest conditions, principally from calcareous, medium-textured glacial drift. The soils are grayish brown or light brown with yellowish-brown subsoils which in the heavy types are comparatively free from coarse material. Areas of Miami soils are usually smooth, with long and gentle slopes. The soils mapped in this series are Miami silt loam with a shallow phase and Miami loam.

The Coloma series includes light-colored upland forest soils developed from sandy materials which were transported through glacial action from sandstone formations principally, and to a less degree from crystalline formations. The mapped types of this series are Coloma fine sand and Coloma sand.

The Warsaw series includes dark-colored prairie soils developed from calcareous, water-laid gravel. These soils occur on stream terraces or outwash plains, and are not subject to inundation. The types mapped in this series are Warsaw silt loam, Warsaw loam, and Warsaw fine sandy loam.

The Fox soils are light colored like the Bellefontaine soils, with parent material of stratified gravel and sand which contains a high proportion of limestone. The land is level and the subsoils are sandy or gravelly at depths ranging from 2 to 4 feet. The soils mapped in this series are Fox silt loam, Fox loam, and Fox sandy loam.

The Plainfield series includes light-colored soils developed from alluvial material derived mostly from glacial débris originating from sandstone rocks. This material is sandy and occurs on stream terraces or outwash plains above the present flood plains of the streams. The mapped soils of this series are Plainfield sand and Plainfield fine sand.

The Superior soils have light-colored surface soils with subsoils of heavy red clay. The parent material was deposited in quiet waters and later modified by glacial action. The land is level. The only Superior soil mapped is fine sandy loam.

Maumee soils are black, poorly drained alluvial soils which have sandy material in the deep subsoil. They are similar to the Warsaw soils except that they are lower and more poorly drained. The subsoils are usually calcareous. Maumee soils mapped are silt loam and loam.

The Clyde series includes dark-colored soils which occur in poorly drained depressions in the upland, and also some lowland soils where the parent material is unsorted glacial till. Clyde silt loam and Clyde loam are mapped in this county.

Poygan soils have dark-colored surface soils with heavy, red clay subsoils developed from heavy-textured, water-laid deposits, under poor drainage conditions. Poygan clay loam and Poygan silt loam are mapped in Green Lake County.

The Dunning series includes dark-colored sandy soils in poorly drained areas. The soils are acid and occur usually outside the limestone region. Only one member of this series, Dunning fine sand, was mapped in this county.

The Genesee series includes light-colored first-bottom land which is subject to flooding. Genesee silt loam is the only soil of this series mapped in this county.

Peat consists of accumulations of vegetable matter in various stages of disintegration and decomposition, mixed with more or less mineral matter. This land is low and very poorly drained.

Rough stony land includes steep, rough, rocky, or broken areas, which may have outcrops of limestone, sandstone, and granitic rocks.



In the following pages of this report the different soils are described in detail and their relation to agriculture discussed; the accompanying soil map shows their location and distribution in the county; and the following table gives their names and extent:

*Acres and extent of Green Lake County soils*

Type of soil	Acres	Per cent	Type of soil	Acres	Per cent
Parr silt loam.....	26,112	13.7	Warsaw fine sandy loam.....	832	0.4
Level phase.....	5,504		Warsaw silt loam.....	512	.2
Parr loam.....	1,536	1.0	Fox sandy loam.....	1,920	.8
Gravelly phase.....	704		Fox loam.....	1,280	.5
Parr fine sandy loam.....	2,176	.9	Fox silt loam.....	448	.2
Dodgeville silt loam.....	1,856	.8	Plainfield sand.....	7,360	3.2
Bellefontaine fine sandy loam, dark-colored phase.....	25,472	11.1	Plainfield fine sand.....	4,480	1.9
Bellefontaine sandy loam, dark- colored phase.....	15,488	6.7	Superior fine sandy loam.....	960	.4
Bellefontaine loam.....	11,200	4.9	Maumee silt loam.....	8,448	3.7
Bellefontaine silt loam.....	6,592	2.9	Maumee loam.....	7,936	3.4
Bellefontaine gravelly loam.....	1,344	.6	Clyde silt loam.....	6,912	3.0
Bellefontaine gravelly sandy loam.....	1,344	.6	Clyde loam.....	1,600	.7
Miami silt loam.....	15,296	7.3	Poygan silt loam.....	2,944	1.3
Shallow phase.....	1,728		Poygan clay loam.....	2,688	1.2
Miami loam.....	3,008	1.3	Dunning fine sand.....	4,416	1.9
Coloma sand.....	7,808	3.4	Genesee silt loam.....	64	.1
Coloma fine sand.....	6,976	3.0	Peat.....	31,552	17.3
Warsaw loam.....	1,344	.6	Shallow phase.....	8,256	
			Rough stony land.....	2,304	1.0
			Total.....	230,400	-----

#### PARR SILT LOAM

Parr silt loam is a prairie soil the topsoil of which consists of very dark brown or nearly black silt loam, rich in organic matter to a depth of 12 or 14 inches. The upper part of the subsoil is dingy-brown or brown silt loam material, which becomes yellowish brown with depth. To a depth of about 3 feet, the lower part of the subsoil is yellowish-brown silty clay loam material. Underlying this is the substratum which consists of very calcareous glacial till composed of clay, silt, sand, and gravel. The line of demarcation between the yellowish-brown silty clay loam material and the substratum is well defined, the upper part being free from boulders and gravel and almost entirely leached of calcium carbonate, whereas the underlying till contains many stones and boulders and much gravel, and effervesces with hydrochloric acid. Both the Truog and litmus-paper tests indicate that the surface soil is acid and needs lime.

Included in mapped areas of this soil are some patches of soils which are not typical Parr silt loam. Where the land is level or almost level, the dark-colored surface layer is thicker, darker, and contains more organic matter than where the surface is rolling. On some steep slopes and small ridges, the unweathered glacial till is nearer the surface than it is in the typical soil. In places where Parr silt loam borders Miami soils it is somewhat lighter in color and contains less organic matter, but where it borders Clyde soils it is darker, contains more organic matter, and its subsoil in some places is mottled. On some gravelly knolls the underlying bedrock of limestone is near the surface, and from these places limestone for agricultural use may be procured.

Parr silt loam is the most important agriculturally, and, with the exception of peat, is the most extensive soil in Green Lake County.

It is the predominating soil of the prairie region which includes some of the most productive soils of the State. Parr silt loam is confined almost entirely to the eastern half of the county, the largest areas being in Green Lake and Mackford Townships where unbroken areas of approximately 20 square miles or more occur. Smaller and more irregular tracts are in Berlin, Brooklyn, and Manchester Townships. In Green Lake and Mackford Townships, Parr silt loam generally adjoins Miami silt loam, but in Berlin and Brooklyn Townships it adjoins Bellefontaine soils.

The land is level or gently rolling and the natural drainage is good except where the surface is level or nearly so. Such areas are mapped separately as Parr silt loam, level phase. The soil as a whole retains moisture well and because of its large supply of organic matter it withstands drought better than most of the other soils of the county.

Parr silt loam was originally covered with a growth consisting chiefly of prairie grasses. Adjoining light-colored soils and in a few other places there were scattered clumps of oak trees and along drainage ways and in other moist places some soft maple. In some places small wild-cherry trees now grow along fence rows and some forest trees have been planted on farmsteads and along some of the highways. More than 95 per cent of this land is improved, almost the only untilled land being that along fence rows and roadsides and around buildings.

The chief crops are corn, hay, oats, barley, wheat, and alfalfa; and hemp and peas are two important special crops. Sugar beets, potatoes, and flax are also grown to a small extent. General farming, including grain growing and dairying is the most important form of agriculture on this land. More beef cattle and sheep are raised on the prairies than elsewhere in the county and many farmers derive a large part of their income from the sale of hogs.

Hemp growing<sup>2</sup> has been developed in Wisconsin within the last 10 years and the center of the industry includes the prairie region of Green Lake County. The production of this crop requires special machinery for harvesting and this machinery is usually owned jointly by the farmers or by the owners of hemp mills where the fiber is prepared for market.

The farms on the prairies are somewhat larger than elsewhere and the system of farming is more extensive. Where dairying is being extended there is a tendency to reduce the size of the farms, but throughout the county as a whole the average size is increasing.

Land values on the prairies are generally higher than elsewhere in the county. Farms on which Parr silt loam predominates range in current value from \$150 to \$250 an acre, depending upon location, roads, and farm improvements. The current value of land without buildings is approximately \$100 an acre.

Analyses and field observations made by the Wisconsin State Soils Laboratory show that Parr silt loam needs lime and phosphate. To meet these requirements 2 or 3 tons of ground limestone per acre should be applied, and about 200 pounds of 16 per cent acid phosphate or one-third as much treble superphosphate per acre. These applications should be used in addition to stable manure. This soil

<sup>2</sup> For a complete discussion of hemp growing in Wisconsin see the following: WRIGHT, A. H. WISCONSIN'S HEMP INDUSTRY. Wis. Agr. Expt. Sta. Bul. 293, 46 p., illus. 1918.

is well supplied with organic matter, but the turning under of vegetable matter will increase its productivity.

*Parr silt loam, level phase.*—The level phase of Parr silt loam is mapped separately because of its topography. It occurs on level or nearly level upland, mostly as patches within areas of the typical soil. It differs from typical Parr silt loam in that the soil is more nearly black, the surface soil is deeper, and there is a tendency toward mottling in the subsoil. It is not so well drained as the typical soil, and in places it requires artificial drainage for the successful production of cultivated crops.

Areas of level Parr silt loam adjoin areas of typical Parr silt loam in the southeastern part of the county, principally in Green Lake and Mackford Townships.

#### PARR LOAM

To depths varying from 12 to 14 inches the topsoil of Parr loam consists of dark-brown or black friable loam. The upper part of the subsoil is rather dark brown loamy material to depths ranging from 18 to 24 inches, where the color is yellowish brown and the texture is slightly heavier. The lower part of the subsoil is loamy material which ranges in texture from fine sandy loam to clay loam. The parent material is gravelly, medium-textured, glacial till in which limestone predominates.

The total area covered by this soil is about 1,500 acres. It is closely associated with Parr silt loam in Berlin, Brooklyn, Mackford, Manchester, and Green Lake Townships. The areas range in size from a few acres to less than half a square mile.

Areas of Parr loam may be level or gently rolling and the drainage is good. The coarse material allows excess moisture to move downward through the soil and be drained away, but sufficient clay is present to retain moisture very well.

This is a prairie soil which originally was covered with prairie grasses. In a few places it was thinly wooded with oak and a few wild cherry trees.

Practically all of the Parr loam is under cultivation and highly improved. About the same crops are grown and the same methods of farming are followed as on the silt loam, but the yields probably average a little lower. The methods of improvement are practically the same as those suggested for Parr silt loam.

*Parr loam, gravelly phase.*—To a depth of 6 or 8 inches the topsoil of Parr loam, gravelly phase, consists of dark-brown or nearly black fine sandy loam, loam, or silt loam. The subsoil is yellowish-brown, gritty silty clay loam material to a depth of about 2 feet, where it is gravelly clay loam material. Crystalline bowlders occur on the surface and rounded limestone gravel throughout the soil. In places beds of gravel occur in the lower depths and in a few places are within 12 inches of the surface. On ridges where the limestone bedrock is not far below the surface, limestone and chert fragments are present on the surface and throughout the soil, and a few outcrops of limestone occur.

Tests made with litmus paper in the field indicate that the surface soil of this phase is acid even where the limestone gravel comes close to the surface. The subsoil, composed principally of material de-



rived from limestone which has not been leached so much as the surface soil, is well supplied with calcium carbonate.

This soil covers about 700 acres, the largest areas occurring in the southeastern part of Mackford Township on low ridges of a narrow belt which extends in an east-west direction. One of these ridges which crosses section 22 is more than a mile long. The ridges usually range in height from 5 to 20 feet and interfere very little with cultivation. Patches of this soil, many of which are too small to be mapped separately, occur in Green Lake, Brooklyn, and Berlin Townships.

Because of the open character of this soil and because the land is uneven, natural drainage is good and in some places excessive. Crops suffer from drought, especially where the gravel comes within 18 inches of the surface. Most of this soil is farmed along with Parr silt loam, but it is not so productive. The crops grown and methods of cultivation are practically the same as on Parr silt loam. Because the subsoil contains much limestone, this soil is better suited to alfalfa than Parr silt loam. The soil warms up early in the spring and may be cultivated soon after heavy rains. If an area is large enough to be made into a field by itself, it is used as permanent pasture.

#### PARR FINE SANDY LOAM

The topsoil of Parr fine sandy loam consists of dark grayish-brown fine sandy loam, varying in depth from 10 to 14 inches. The subsoil material is stiff, yellowish-brown fine sandy clay loam underlain at depths of about 2 or 3 feet by yellowish silty clay loam material. Mapped as Parr fine sandy loam are some areas which differ from the typical soil in that the color of the surface soil is light gray owing to lack of organic matter; some small areas where the surface soil is darker than typical; some areas where the underlying formation is within 3 feet of the surface and the subsoil material is sandy, as in section 20, Kingston Township; and some patches where the surface soil is very fine sandy loam.

Parr fine sandy loam covers a total area of 3.4 square miles. It occurs principally in the southern half of the county, the largest area being mapped in the central part of Kingston Township, and others in Manchester and Mackford Townships.

The surface of Parr fine sandy loam is mostly gently rolling with a few steep slopes, but modern machinery can be used to advantage in cultivation. Because of the surface features and the sandy nature of the soil the natural drainage is good. The subsoil is sufficiently heavy in most places to be retentive of moisture.

Parr fine sandy loam is good agricultural soil, although not so productive as Parr silt loam. It warms up early in spring and responds readily to fertilization and improved methods of cultivation. Although this soil is used chiefly for general farming and dairying, it is naturally better suited to truck crops. Corn does very well because it may be planted comparatively early in the spring. More rye is grown on this soil than on the heavier soils. It is also better adapted to the production of potatoes than is Parr silt loam. On account of the acid condition of the soil, some diffi-

culty is experienced in maintaining a stand of clover or alfalfa, but by the use of lime these crops can be grown. The same crops are grown and the same methods of culture followed as on Parr loam and Parr silt loam, but Parr fine sandy loam has a somewhat lower value. In the improvement of this soil the acid condition should be corrected and the need for organic matter and phosphorus supplied.

#### DODGEVILLE SILT LOAM

To a depth of 10 or 12 inches, the surface soil of Dodgeville silt loam consists of dark-brown or nearly black silt loam. This is underlain by a rather dark brown silt loam layer which gradually becomes yellowish brown in color and heavier in texture with depth. The subsoil below a depth of 24 inches is yellow or yellowish-brown silty clay loam or clay loam material which contains sufficient fine gravel, grit, and small stones to make it decidedly gritty. At depths ranging from 2 to 3 feet is the substratum of unsorted glacial till which rests upon limestone bedrock.

On some areas, the gritty subsoil is within 12 or 18 inches of the surface, and in a few places the underlying rock is within 2 feet of the surface. The color of Dodgeville silt loam is usually lighter than that of Parr silt loam, and the surface is more irregular. Gravel, crystalline boulders, and angular limestone fragments occur on the surface and throughout the soil.

Dodgeville silt loam occurs throughout the prairie region where it is associated with Parr silt loam. It is inextensive, covering only about 5 per cent of the prairie region, and occurs as long, narrow strips along the tops of low ridges or on the steeper slopes of small hills and ridges. Because the land is sloping and the subsoil gravelly, drainage is everywhere good and in a few places excessive.

Dodgeville silt loam requires the same treatment as Parr silt loam, but it is less acid because the limestone material is closer to the surface, and it is consequently better adapted to growing alfalfa. As drainage is better than on Parr silt loam, the soil can be worked a little earlier in spring and sooner after rains, but as this soil occurs in such small patches advantage is seldom taken of this condition. The same crops are grown and the same methods of farming are followed as on Parr silt loam, but the soil is not considered so valuable.

#### BELLEFONTAINE FINE SANDY LOAM, DARK-COLORED PHASE

The topsoil of Bellefontaine fine sandy loam, dark-colored phase, consists of friable, dark-brown fine sandy loam to depths varying from 8 to 12 inches. This is underlain by reddish-brown, calcareous, gravelly loam or gravelly clay loam material, which continues to a depth of 2 or 3 feet where it grades into unsorted glacial till consisting of a mixture of sand, clay, gravel, and boulders.

In Brooklyn Township there are some areas on which the surface soil is very fine sandy loam. In areas near the prairies, the color of the soil is darker than typical. Where the gravelly layer is near the surface the soil approaches a gravelly loam in texture. In many places some gravel is present on the surface as well as in the soil, but where this condition exists extensively such areas are mapped as Bellefontaine gravelly loam or Bellefontaine gravelly sandy loam.



There are some boulders on the surface and in some places these are so numerous that they have been gathered from the fields and piled into stone fences. In some areas the color of the subsoil is yellowish brown instead of reddish brown as in the typical soil and the material is less gritty. Adjoining areas of Coloma soils, the surface of this soil is more sandy and on a few small patches the fine sand layer extends to a depth of 2 feet.

Bellefontaine fine sandy loam, dark-colored phase, is one of the important and extensive soils of the county. There are no large unbroken tracts, but there are a large number of areas where the continuity is broken by patches of other soils of the Bellefontaine series and of the Miami series. This soil occurs throughout the county, and is most extensively developed in Brooklyn, Manchester, Princeton, Marquette, and Kingston Townships.

The land for the most part is gently rolling, although in some places it is nearly level and in others irregular. Only where the slopes are steep or the ridges are sharp is the use of modern farm machinery impracticable. The level areas are inextensive, but are the most desirable. Drainage is good, but there is sufficient clay in the subsoil to retain moisture, and only the more sandy areas suffer from drought.

In the remaining wood lots oak is now the chief growth. Some fair timber still remains, but most of the merchantable timber has been cut. Probably 80 per cent of this soil is farmed. It is much esteemed because of its easy tillage and quick response to fertilization. It is fairly productive. The acreage in corn is about 30 per cent greater than that in oats or mixed clover and timothy, the crops next in importance. Rye, barley, potatoes, alfalfa, wheat, and melons are also grown. There are many small apple orchards, but the fruit is usually of inferior quality.

The soil is treated in practically the same way as the Miami and Bellefontaine loams. Corn is followed by small grain, and this by mixed timothy and clover. Very little fertilizer other than barnyard manure is used, although marked increases in yields of almost all crops have been obtained by the use of acid phosphate. With an application of ground limestone, a much better stand of clover and alfalfa is obtained.

Although Bellefontaine fine sandy loam, dark-colored phase, is easily cultivated and improved, a decrease in the productiveness of many farms has taken place, owing to neglect to rotate crops, grow clover, and prevent the washing away of the soil on the hillsides. The steeper slopes should be seeded to grass and used for permanent pasture, or for growing alfalfa. On the more gentle slopes contour cultivation and the growing of winter crops of rye or wheat following corn is advisable. The suggestions offered for increasing the content of organic matter and for correcting acidity in Miami silt loam and Parr silt loam are applicable to this soil. Potatoes, beans, cucumbers, strawberries, and tomatoes grow well, and their cultivation might profitably be extended in locations favorable for marketing.

#### BELLEFONTAINE SANDY LOAM, DARK-COLORED PHASE

The surface soil of Bellefontaine sandy loam, dark-colored phase, consists of dark grayish-brown sandy loam to a depth of 6 or 8

inches. This is underlain by a 10-inch layer of yellowish-brown sandy loam, which in turn is underlain by reddish sandy clay loam material, which becomes somewhat heavier at a depth of about 2 feet, and is in places stiff reddish silty clay loam. The typical unsorted calcareous glacial till consisting of sand, limestone, gravel, and some clay occurs from 30 to 36 inches below the surface.

This soil varies in texture from sand to loam. The topsoil may consist of loamy medium sand, but where this is the case, the deep subsoil usually contains enough clay to give the soil the agricultural value of sandy loam. Where sandstone lies near the surface, the subsoil is sand. In section 20, Kingston Township, and in a few other areas, the material rests on a sandstone formation at depths of less than 3 feet. On some knolls and slopes gravel appears on the surface and may be mixed with the soil. However, gravel, stones, and boulders are not so numerous on this soil as on Bellefontaine loam and Bellefontaine fine sandy loam, dark-colored phase. The supply of organic matter in the surface soil is uniformly low, and most of the surface soil is acid.

This soil is most extensively developed in the townships of Kingston, Marquette, Berlin, Princeton, and Manchester, the largest area covering 5 or 6 square miles in the central part of Berlin Township. Bellefontaine sandy loam, dark-colored phase, is closely associated with Bellefontaine fine sandy loam, dark-colored phase, and Bellefontaine loam, and in many places borders Coloma fine sand.

The surface of this soil is mostly gently rolling and modern farm machinery can be used on most of it. However, in western Kingston Township and at a few other places steep slopes occur, where there is danger from erosion and where self-binders and tractors can not be used to advantage. Where sandstone lies within 3 or 4 feet of the surface the land is level or very gently sloping and in places has the appearance of terrace land. Because the surface is undulating and the soil is gravelly and sandy, natural drainage is good, and in the more sandy areas is excessive.

Some of the steep slopes are forested, chiefly with oaks and hickory, but the greater proportion of the soil is cultivated and is in improved farms. Being sandy, it is not so desirable as Bellefontaine loam or silt loam, but it is fair agricultural land, and many good farms are located on it. It has the advantage of warming up early in the spring, is easily cultivated, well drained, and responds quickly to fertilization.

The chief crops are corn, oats, rye, barley, potatoes, and wheat. Hay is also grown, but not in such quantities as on the heavier soils. There are good fields of alfalfa, especially where the gravelly clay loam is close to the surface and the soil is least in need of lime. Rye is grown more extensively than on the heavier soils and gives good yields. Some of the farmers reported very good yields of corn. In Berlin Township, where large areas of this soil occur, there was in 1922 an average of 18 acres of corn and 23 acres of oats on each farm. In Kingston Township, where large areas of this soil are under cultivation, there was an average of 24 acres of corn and 14 acres of oats on each farm. In Mackford Township, where Parr soils predominate, there was an average of 27 acres of corn and 30 acres of oats on each farm. These varying crop acreages are the

result of differences in the character of the soils. The same methods of cultivation, fertilization, and crop rotation are followed on this soil as on Bellefontaine loam and fine sandy loam, dark-colored phase.

In the improvement of this soil the organic-matter content should be increased by growing legumes more extensively; phosphorus should be supplied by using acid phosphate to supplement the stable manure or the green-manure crop, and ground limestone should be used where the soil is acid, especially where there has been difficulty in obtaining or maintaining a stand of clover or alfalfa.

#### BELLEFONTAINE LOAM

The surface soil of Bellefontaine loam, to depths of 8 or 10 inches, is brown loam which usually contains some gravel. This is underlain by reddish-brown clay loam material to depths ranging from 20 to 30 inches. Below this lies a shallow layer of gravelly clay loam or loam material, underlain by stony, calcareous, gravelly till composed largely of limestone. In some places this layer of till is so close to the surface that the surface soil approaches a gravelly loam or gravelly sandy loam in texture. A few crystalline boulders are present on the surface and throughout the soil and substratum. In section 19, Manchester Township, and in a few other places, where the surface is almost level, areas of this soil are underlain by bedded limestone at depths between 30 and 40 inches. There are other level tracts where the surface soil is darker than typical and contains more organic matter. In the western part of the county are patches in which the deeper subsoil material consists of sand, owing to its proximity to underlying sandstone.

Areas of Bellefontaine loam are well distributed throughout Green Lake County, the most extensive occurring in Manchester Township, and smaller tracts in Marquette, Kingston, Green Lake, Brooklyn, Princeton, and Berlin Townships. This land is undulating or very choppy, with numerous knolls, ridges, and depressions. Because of this irregular surface and the open gravelly subsoil, the natural drainage is good.

A large proportion of Bellefontaine loam is under cultivation, but some of the land is forested with oak, hickory, and maple. All crops common to this region grow well on this soil. Corn, oats, and hay are the principal crops. Alfalfa and clover do very well. Dairying is carried on extensively. The manurial requirements and crop adaptations of this soil are similar to those of Bellefontaine silt loam, and the suggestions offered for the improvement of Bellefontaine silt loam apply also to this soil.

#### BELLEFONTAINE SILT LOAM

To depths ranging from 6 to 8 inches, the cultivated surface soil of Bellefontaine silt loam is mellow brown or grayish-brown silt loam low in organic matter. Below this, to depths varying from 12 to 20 inches, is a layer of yellowish-brown or slightly reddish brown silty clay loam. The subsoil of reddish-brown silty clay loam material is plastic and sticky when moist and hard when dry, and continues to depths ranging from 24 to 30 inches. Below this is a shallow layer of friable, gravelly, moderately calcareous loamy material, underlain by the parent material which is a heterogeneous mass of



unweathered gravelly till originating principally from limestone. The surface soil is slightly gravelly, and the subsoil contains enough sand to make it gritty. Some crystalline bowlders occur on the surface and throughout the soil and subsoil. In a few areas near the prairies the color of the surface soil is darker than typical. In section 24, Kingston Township, where the slopes are steep, the limestone substratum lies near the surface.

Areas of Bellefontaine silt loam, as mapped, include numerous small patches of other soils, such as Bellefontaine gravelly loam, loam, fine sandy loam, and Miami silt loam. Sandy and gravelly areas usually occur on knolls and steep slopes, whereas the more silty areas are gently undulating and subjected to very little erosion. Bellefontaine silt loam differs from Miami silt loam in that the brown surface soil is darker, the subsoil has a reddish tinge, and the calcareous parent material is nearer the surface.

The largest areas of Bellefontaine silt loam are in Brooklyn Township, and smaller areas are in Berlin, Green Lake, Manchester, Kingston, and Mackford Townships. The land is moderately undulating to choppy and broken, but most of it is smooth enough for the use of improved machinery in planting, cultivating, and harvesting crops. Surface and internal drainage are good.

It is estimated that 75 per cent or more of this soil is under cultivation. The principal crops are corn, oats, and mixed clover and timothy with a considerable acreage of rye, barley, alfalfa, potatoes, beans, and wheat. Nearly every farm has a few apple trees, but there are no important commercial orchards. The yields of all crops are equal to or somewhat greater than those grown on Miami silt loam. In the improvement of this soil the supply of organic matter and nitrogen should be increased. This may be done by increasing the legume crops. The use of stable manure may be supplemented by the use of acid phosphate as the soil is somewhat deficient in phosphorus.

#### BELLEFONTAINE GRAVELLY LOAM

The topsoil of Bellefontaine gravelly loam is light-brown or brown loam, carrying considerable gravel and is from 6 to 10 inches deep. The subsoil is yellowish-brown or reddish-brown, friable, gravelly loam or gravelly clay loam material to depths ranging from 30 to 40 inches. Below this is the parent material of coarse, calcareous glacial till. Many small patches of Bellefontaine gravelly fine sandy loam, fine sandy loam, loam, and silt loam are included with mapped areas of this soil. In places inclusions of silt loam are so numerous that it is difficult to determine whether the silt loam or the gravelly loam predominates. Beds of gravel occur in many places in the deep subsoil.

Areas of this soil, although inextensive, are widely distributed throughout the county, occurring in Manchester, Marquette, Berlin, Brooklyn, St. Marie, and Princeton Townships. Many patches too small to indicate separately are mapped with such soils as Bellefontaine sandy loam, dark-colored phase, fine sandy loam, dark-colored phase, and loam. This soil occurs on small knolls, hills, and ridges, or on slopes bordering drainage ways. On most of this land modern farm machinery can be used, but slopes on which it would be difficult or impracticable to use a tractor or self-binder, are numerous,

Because of the unevenness of the surface and the gravelly surface soil and subsoil, natural drainage is good and in many places excessive. Where the land is level this soil is well adapted to alfalfa and clover, probably more so than any other soil in the county, chiefly because of its abundant supply of lime.

The crops grown, methods used, and yields obtained compare very favorably with those on Bellefontaine loam. A larger proportion of this soil is in wood lots and permanent pasture than is the case with other upland soils. The rough areas afford good grazing and the land can be utilized to good advantage where dairying is carried on.

#### BELLEFONTAINE GRAVELLY SANDY LOAM

Bellefontaine gravelly sandy loam is, for the most part, brown sandy loam containing gravel, underlain by reddish gravelly clay loam material, which in places is heavy. The subsoil in most places is underlain by beds of gravel, or extremely gravelly glacial till, principally limestone. The surface soil is not acid or only slightly so, probably owing to the influence of the lime from the gravel. Some glacial bowlders are scattered over the surface and throughout the soil. In some areas the soil is unusually sandy, but in others the surface soil is loam.

Areas of Bellefontaine gravelly sandy loam are inextensive although widely distributed throughout the county, occurring on small knolls or narrow ridges in Mackford, Manchester, Marquette, Princeton, and Berlin Townships. The land varies from gently rolling to choppy. Because of the uneven surface and gravelly subsoil, natural drainage is good and in places excessive.

The smoother areas of Bellefontaine gravelly sandy loam are cultivated in the same manner and planted with the same crops as Bellefontaine sandy loam, dark-colored phase, whereas the steep slopes and ridges are, for the most part, either forested or used for permanent pasture. Because of its sweetness, this soil is well adapted to alfalfa. It is somewhat lower in agricultural value than the heavier Bellefontaine soils. Suggestions offered for the improvement of Bellefontaine sandy loam, dark-colored phase, Bellefontaine fine sandy loam, dark-colored phase, and Bellefontaine loam, apply also to this soil.

#### MIAMI SILT LOAM

The cultivated surface soil of Miami silt loam is a layer from 8 to 12 inches thick, of light grayish-brown, smooth silt loam comparatively low in organic matter. This is underlain by yellowish-brown, heavy silt loam which gradually becomes heavier with depth changing to silty clay loam, and to yellow clay or clay loam below a depth of 20 or 24 inches. This heavy material continues to a depth of 3 or 4 feet, where gritty or gravelly calcareous material is mixed with the clay and silt. The soil within 3 feet of the surface is loesslike in structure and texture and free from coarse material; but the deeper part of the subsoil is chiefly unassorted till consisting of calcareous clay, sand, gravel, and fragments of limestone and crystalline rock.

Miami silt loam as mapped in Green Lake County differs from the Miami silt loam that occurs in most of Ohio, Indiana, and southern

Michigan, in having a greater average thickness of surface silty soil, and a greater total thickness of weathered material leached of calcium carbonate.

A few boulders are present in some areas, but as a whole the land is free from them. The underlying rock is chiefly limestone, and this crops out on some of the steeper slopes. The surface soil, according to the litmus-paper and Truog tests is slightly acid. This soil is very uniform in color, but where it borders the Parr soils of the prairies it is darker than the typical soil and contains more organic matter. In such places the change is so gradual from one kind of soil to the other that no definite boundary can be determined.

Miami silt loam differs from Bellefontaine silt loam in that it has a more pronounced gray color, a smaller quantity of coarse material, a greater thickness of silty surface soil, and fewer surface stones and boulders. Both are light-colored upland forested soils and are spoken of as "clay land" to distinguish them from the darker-colored prairie soils.

The total area of Miami silt loam is approximately 24 square miles. It usually borders the prairies, the largest areas occurring in the townships of Mackford and Green Lake, and small patches in the townships of Manchester, Brooklyn, Berlin, and Marquette.

This land is gently rolling, and modern farm machinery can be used on more than 90 per cent of it. The most elevated surfaces of the county are covered by prairies. On the slopes extending from these Miami silt loam has been developed. Adjacent to the prairies are level areas where drainage is poor and the surface soils are darker and deeper than typical Miami silt loam. Farther down on these slopes and also on slopes bordering Green Lake and various drainage ways are long narrow belts with comparatively steep rough surfaces where there is danger of erosion. On such surfaces the soils are well drained and withstand droughts better than most of the soils of the county.

Miami silt loam is one of the most productive soils of the county, and approximately 80 per cent of it is under cultivation. Some of the areas along the south shore of Green Lake and on the steeper slopes along Grand River are forested and in places contain considerable merchantable timber. The growth consists chiefly of several varieties of oak, elm, hickory, maple, basswood, and wild cherry.

General farming, with dairying, is the principal form of agriculture. The most important crops are corn, hay, oats, barley, wheat, with some peas, alfalfa, and a little hemp. Potatoes are grown chiefly for home use.

This soil is treated in about the same manner as Parr silt loam, but because it contains more lime it is better adapted to clover and alfalfa. Less corn and oats are grown on this soil than on Parr silt loam.

In the cultivation of this soil fall plowing is common, although on account of its friability it can be worked early in the spring. The subsoil is retentive of moisture, and a good surface mulch enables it to resist drought. Barnyard manure is the principal fertilizer, and as dairying is extensively practiced a good supply is always available. From 10 to 15 loads an acre are usually applied every four or five years to sod land to be plowed for corn. Green-manure crops are used to a small extent. Cultivation is not usually difficult as the



surface soil is very silty and pulverizes readily, but where the subsoil is exposed on slopes or where the surface soil is shallow and the heavier material is turned up by the plow, some difficulty is experienced.

The usual crop rotation consists of corn, barley, and oats, one year each, followed by mixed timothy and clover. Wheat may take the place of oats or barley, or it may be made another step in the rotation. Grass is usually left for two years, being cut for hay the first year, and used as pasture the second year. In localities where there is permanent pasture land, as on areas of Clyde soils and peat, the upland soil is used very little for pasture.

One of the special crops on this soil is sugar beets. Yields average from 12 to 15 tons an acre, and the sugar content is high. Because the soil is silty and friable, it is comparatively easy to maintain a good surface mulch, and the work of cultivating the crop is less arduous than on heavier soils. Alfalfa is considered an important crop, but its acreage does not equal that of clover. Most farmers grow potatoes for home use and frequently have some to sell. Normal yields range from 150 to 200 bushels an acre and the quality is good. Hemp is also an important though a rather new crop grown on this soil.

The most important factor in cultivating Miami silt loam is the need of increasing the organic matter, which can best be accomplished by liberal applications of manure, but as the supply of stable manure is inadequate to meet the needs of the entire farm, green-manure crops should be grown and plowed under. The acid condition of the soil may be corrected by the application of ground limestone, which has a tendency to increase the yields by making soil conditions more favorable for plant growth.

Stable manure treated with acid phosphate may be used to overcome any lack of phosphorus. If the supply of manure is insufficient and legumes are not grown for green manure, mixed commercial fertilizers should be used; but the use of legumes as green manure is preferable to the use of commercial fertilizers. Where trucking is carried on or where special crops are grown, mixed fertilizers may be used.

Dairying could be brought to a higher stage of development on this soil. More alfalfa should be grown and every dairy farmer should include it in his farm crops. More hogs could also be raised with profit.

*Miami silt loam, shallow phase.*—Miami silt loam, shallow phase, occurs on ridges, slopes, and hilltops, where at depths ranging from 12 to 30 inches the soil material is underlain by limestone or unsorted glacial till. On such areas the topsoil is practically the same as that of typical Miami silt loam, but the subsoil contains more gravelly material, and drainage is better. In general, soil of this phase is cultivated in about the same way as typical Miami silt loam.

#### MIAMI LOAM

The cultivated topsoil of Miami loam consists of a 10-inch layer of light-brown loam, underlain to a depth of about 20 inches by yellow or yellowish-brown loam. The subsoil is friable silty clay loam material to depths of about 3 feet, where it is underlain by

parent material of unassorted glacial till consisting of sand, gravel, and clay. To a depth of 3 feet the soil is free from coarse material, and in this respect it differs from Bellefontaine loam.

Mapped areas of Miami loam include some patches of very fine sandy loam, in Brooklyn Township, some small areas where the subsoil is very fine sand, and others where the soil is dark brown and heavier than in the typical soil. Areas of Miami loam are inextensive, occurring in the townships of Manchester, Green Lake, Brooklyn, and Berlin. As a rule, this land is gently sloping and is well suited to the use of modern farm machinery. Drainage is good, except in a few places where the land is level.

This soil was originally forested, principally with oak and hickory, but most of it is now cleared and used for general farming and dairying. The chief crops are corn, hay, small grain, and alfalfa. The same methods of cultivation and fertilization are used as on Miami silt loam, resulting in about the same average yields. This soil is deficient in organic matter, nitrogen, phosphorus, and lime. More legumes should be grown to increase the supply of organic matter. The application of acid phosphate would insure larger yields of all crops.

#### COLOMA SAND

Coloma sand consists of light-brown fine or medium sand to a depth of 6 inches, where it is underlain by yellow fine sand, in places containing some fine gravel. The topsoil and subsoil are loose and open and contain but little material as fine as silt. In some areas the subsoil is sticky.

The larger areas of Coloma sand are in the vicinity of Berlin and in the northwestern part of the county. The surface is gently rolling, and in no place is the land too steep for cultivation. The drainage is very good. The native growth on this soil consists chiefly of scrub oak and jack pine. Heath growths have come up on abandoned fields. Some fields have remained idle for several years because of the small yields secured. Where the soil is cultivated, the chief crops are rye and potatoes, less important crops being hay and beans, but yields are small and unsatisfactory. Where crops are forced to mature early and where it is profitable to use large quantities of soluble fertilizers, this soil is especially suitable for truck crops. For the improvement of this soil, legumes should be grown. Small grain, clover, followed by corn is a good three-year rotation, in which the second cutting of clover should be plowed under.

#### COLOMA FINE SAND

Coloma fine sand is similar to Coloma sand, differing mainly in the texture of the surface soil. Also the surface soil contains more organic matter, and its moisture-retaining capacity is greater.

Mapped as Coloma fine sand are some areas where the soil is of different texture, as in western Brooklyn Township where the topsoil and subsoil of some areas consist of very fine sand, and in some other areas silt or clay layers near the surface influence the moisture-retaining capacity of the soil.

Areas of Coloma fine sand occur principally throughout the central part of the county and smaller and more scattered areas in the



northeastern and southwestern parts. Drainage conditions and agricultural possibilities of Coloma fine sand are similar to those of Coloma sand.

#### WARSAW LOAM

Warsaw loam consists of very dark brown or nearly black loam, underlain at a depth of about 10 inches by rather dark brown loam which with increasing depth becomes lighter in color and more sandy in texture. At depths varying from 18 to 24 inches is clay loam material which is underlain by sandy, gravelly material at a depth of about 3 feet.

The total area of Warsaw loam is 2.1 square miles, the largest areas occurring in the southern part of the county along Grand River in Manchester Township.

This land is level or gently sloping and usually borders low, poorly drained soils such as those of the Clyde series. Drainage is usually good, but in some low places it is slightly deficient in the spring when the water table is highest. Where the gravel subsoil is close to the surface, the land may suffer from lack of moisture during dry periods.

This soil has been developed under prairie conditions, and the original growth consisted chiefly of prairie grasses. Some trees bordered the watercourses, but the greater part of the land was treeless.

Practically all this soil is now under cultivation, and is considered fertile and productive. Corn is the chief crop and oats are probably second in acreage. Most of the general farm crops common to the region are grown, and give fairly good returns. The suggestions offered for the improvement of Parr silt loam are applicable also to Warsaw loam.

#### WARSAW FINE SANDY LOAM

The surface soil of Warsaw fine sandy loam, to a depth of 10 inches, consists of dark-brown or nearly black fine sandy loam. This is underlain by material of practically the same texture but of chocolate-brown color. Below 18 inches, a yellowish color appears, and at depths varying from 20 to 28 inches the content of gravel increases very much, and the material grades downward to beds of stratified sand, or sand and gravel. This soil is somewhat variable and in a few places the subsoil is heavier than in the typical soil. The surface soil gives an acid reaction.

Warsaw fine sandy loam is of very small extent and is confined chiefly to the southern part of the county along the valley of Grand River, one of the larger areas being located in section 7 and another in sections 11 and 14 of Kingston Township.

The surface of this soil is level or nearly so, and because of the open subsoil the natural drainage is good.

Warsaw fine sandy loam is an alluvial soil, having been deposited by streams when the volume of water was much greater than at present. The parent material was glaciated material which came in part from limestone and in part from sandstone. This is a prairie soil. Although some trees formerly grew along watercourses, the land as a whole was nearly treeless and the chief growth consisted of prairie grasses.

Warsaw fine sandy loam is a fair agricultural soil. It is all farmed and produces most of the crops common to the region. It is farmed in the same manner as the loam but is better suited to truck crops than to general farming operations. Improvements should be made along the same lines as those suggested for Parr loam and Parr silt loam.

#### WARSAW SILT LOAM

Warsaw silt loam, to a depth of about 12 inches, is very dark brown silt loam, containing a comparatively high proportion of organic matter and some fine sand. The upper subsoil is dark-brown silt loam, grading downward to brownish-yellow silt loam or silty clay loam which contains a small quantity of sand. At depths ranging from 24 to 36 inches, yellow fine sandy loam or sandy loam is encountered which is abruptly underlain by gravelly sandy loam, and this in turn by stratified beds of gravel and sand. The soil, as mapped, includes patches of Warsaw loam not sufficiently extensive to warrant separate mapping. Small areas of a deep phase have also been included with the mapped Warsaw loam. In such areas the stratified beds of sand and gravel are more than 3 feet beneath the surface. In a few places the stratified beds of sand and gravel are within less than 2 feet of the surface.

Warsaw silt loam is developed chiefly in the southern part of the county within the drainage basin of Grand River. Most of this soil is in sections 23 and 24 of Manchester Township and in sections 7 and 8 of Mackford Township, partly within the city limits of Markesan.

The surface varies from level to very gently undulating, but in section 23, Manchester Township, it is rolling. Natural drainage is good, and where sand and gravel beds are within 2 feet of the surface the soil dries out somewhat during dry periods. Warsaw silt loam was originally prairie land and the native growth consisted of prairie grasses. A few trees grew along streams but the land in general was treeless.

Warsaw silt loam is all in well-improved farms, is excellent farming land, and is used for general farming and dairying. Corn, oats, barley, and hay are the principal crops, corn occupying the largest acreage. Some wheat is also grown.

Because areas of this soil are so small, few if any farms are located entirely upon it, but it ranks among the higher-priced farm lands of the county. Improvements suggested for Parr silt loam are applicable to this soil also.

#### FOX SANDY LOAM

The topsoil of Fox sandy loam consists of brown or grayish-brown sandy loam, underlain at depths ranging from 8 to 12 inches by reddish-brown sandy loam. The subsoil below a depth of 20 inches is gravelly sand material, which grades into beds of stratified sand and gravel, the gravel being largely limestone. In some places no gravel is present in the subsoil, and here the subsoil is a yellow sand or fine sand which in places has a reddish cast. Gravel occurs on the surface in places.

Areas of this soil are inextensive, and occur chiefly on the terraces along Grand River and its tributaries in Kingston and Manchester

Townships. Mapped with Fox sandy loam are some small areas where gravel occurs in the surface soil. The largest of these areas is in section 16, Manchester Township.

This land is level or gently sloping, except in a few places where the terraces have been eroded. The soil is open and porous, and drainage is good and in places excessive. Most of the terraces are well above the present flood plains of the streams and none of the soil is subject to overflow.

Most of this land is included in farms, but it is not all well improved. As compared with heavier soils it is of rather low agricultural value, although under careful management yields are fair. Rye and potatoes are grown, and these crops are better suited to this soil than most of the other crops. Corn does fairly well especially if improved methods of fertilization and cultivation are used. It is not good pasture soil, although some hay is grown on it. It is better suited to truck crops than to general farming, but the small size of the areas and their location make truck growing on a large scale impracticable.

In the improvement of this soil the same methods may be followed as are suggested for sand soils such as Plainfield sand and fine sand.

#### FOX LOAM

Fox loam consists of light chocolate-brown friable loam to a depth of 8 or 10 inches where it grades to friable reddish-brown loam. Below a depth of 18 inches a reddish-brown gritty clay loam material occurs, which at a depth of about 2 feet is underlain by stratified, calcareous sand and gravel. Fox loam is very variable in texture, ranging between the heavier silt loams and the more sandy soils. Litmus-paper tests show that the soil is acid in many places, but the deeper part of the subsoil contains much limestone gravel and is neutral or slightly alkaline.

This soil is of small extent, and is developed chiefly in sections 12, 14, 15, 16, and 33 of Manchester Township. Small scattered areas occur in a few other places in the southern part of the county, some of which are too small to be shown on the map. This land is level or very gently sloping, and the drainage is good. The area in section 33, Manchester Township, is about 40 feet higher than the marshland adjoining and from a distance it appears to be a high plain or terrace, but a closer view shows that owing to erosion the surface is somewhat uneven.

Nearly all this soil is under cultivation, and the principal crops are corn, oats, barley, rye, and hay. Some potatoes are grown and the soil is well adapted to truck farming, but this industry is not carried on extensively. The farm practices followed and the yields secured are about the same as those on Fox silt loam. The soil is easy to cultivate, and a good mellow seed bed can be readily developed. The soil is deficient in organic matter. Methods of improvement suggested for Bellefontaine silt loam and other light-colored heavy upland soils also apply to this soil.

#### FOX SILT LOAM

The surface soil of Fox silt loam is medium-brown or dark-brown friable silt loam from 6 to 8 inches deep. This is underlain by



reddish-brown loam or fine sandy loam to a depth of about 20 inches, where fine sand and gravel appear. Mapped areas include numerous patches which are not typical, some of which closely resemble loam, and others are heavier than silt loam. There are patches where the dark-brown friable silt loam material rests on unweathered sand and gravel at a depth of about 3 feet. In a few small areas Fox silt loam is almost as dark as Warsaw silt loam. The surface soil is slightly acid, but the deep, gravelly subsoil is well supplied with lime, the gravel consisting principally of limestone.

The total area of Fox silt loam is less than 1 square mile. Areas are mapped in section 12, Manchester Township, in sections 7, 9, and 10, Mackford Township, and section 35, Green Lake Township.

The land is level or nearly so, except where this soil occurs on steep terrace slopes bordering watercourses. Because of the open subsoil natural drainage is good and in places excessive. Most of this soil is in farms, but because of its occurrence in small areas very few if any farms are made up entirely of it. It is devoted to general farming and dairying. Where the surface is sloping or steep the land is best suited to permanent pasture. Fox silt loam is similar to Bellefontaine silt loam and methods of improving it are the same. The soil is deficient in organic matter and phosphorus and in places it needs lime. Steep slopes should be kept in grass to prevent the formation of gullies and the washing away of the surface soil.

#### PLAINFIELD SAND

Cultivated Plainfield sand to a depth of 8 inches consists of loose, open brown sand containing a small quantity of organic matter. It has a little gravel scattered over the surface and mixed with the soil. The subsoil is yellow medium sand, which becomes coarser in texture with increased depth, and below 30 inches contains considerable coarse sand and fine gravel. In some places where the subsoil contains considerable iron the material has a brownish or a reddish color; where the soil borders marshland the color is darker than typical owing to a larger percentage of organic matter; and where it is adjacent to heavier soils the texture approaches sandy loam and the soil is of better quality. In a few places sufficient clay is present in the subsoil for the material to feel slightly sticky, which increases the agricultural value of the soil. Virgin soil of this kind contains more organic matter within the first inch or two of the surface than is found in old fields, but under cultivation this is soon lost.

Plainfield sand covers a total area of 11.5 square miles, and occurs chiefly in the northern and western parts of the county, the largest areas being in the northern parts of Berlin and Seneca Townships, and many smaller areas in St. Marie, Princeton, and Marquette Townships. Plainfield sand occurs on the terraces along the old flood plain of Fox River.

The surface is level or slopes very gently toward the stream along which the soil occurs. In places it occurs as islands in the marshland country, where it is usually only from 1 to 3 feet higher than the wet land, and here the water table is closer to the surface than typical and the soil may be wet in the spring. Most of this land is high enough to be well drained, and in some places drainage is excessive.

The original forest growth consisted mainly of red and black oak, jack pine, and some Norway pine. The best timber has been cut, but a large number of small jack pine and some oak still remain, the largest of which range from 8 to 12 inches in diameter.

Probably 40 per cent of this land has been cleared and cultivated, but some farms have been abandoned. The chief crops grown at the present time and the average yields during the most favorable years are as follows: Corn, from 15 to 25 bushels an acre; oats, from 20 to 25 bushels; rye, from 12 to 20 bushels; buckwheat, from 12 to 16 bushels; and potatoes, from 100 to 150 bushels. Potatoes are the chief cash crop. A little hay, consisting of clover and timothy, is produced, but the stand is thin and the yield seldom amounts to 1 ton an acre. Some sorgho (sweet sorghum) is grown and yields about 24 gallons of sirup to the acre.

Because Plainfield sand is easily cleared and cultivated, some farmers believe that it is preferable to heavier, higher-priced land. The methods of cultivation, crop rotation, and fertilization followed on many farms are not best suited to soil of this character. Very little livestock is kept, and consequently very little manure is available. Because of this and failure to plow under green-manure crops, the organic matter in the soil is rapidly depleted. The best farmers make a good living from their farms, but most of the farm buildings reflect the generally poor condition of the soil.

The limiting factors in crop production on this soil are low water supply, low fertility, and low organic-matter content. Improvement of this soil must be based on an increase of organic matter, which will not only increase its water-holding capacity and fertility, but will also tend to prevent blowing. This can be done by the growth of legumes and the plowing under of green-manure crops. A rotation consisting of corn, rye, and clover is well suited to sandy soil, and in this rotation the second crop of clover should be plowed under. Mixed fertilizers may be needed in addition to this, but clover will supply most of the much-needed nitrogen and organic matter. Owing to the acid condition of this land, it is necessary to use lime to insure a good growth of clover. Where stock is kept, the manure should be carefully conserved and applied to the land with as little loss as possible. Corn can be grown more profitably on this light sandy soil than potatoes, so corn rather than potatoes is suggested in the rotation.

#### PLAINFIELD FINE SAND

To a depth of 6 or 8 inches, cultivated Plainfield fine sand consists of light-brown or grayish-brown fine sand which contains some organic matter. The subsoil is yellow, fine sandy material to depths of more than 3 feet. Mapped areas of Plainfield fine sand include variations from the typical soil in which the subsoil material is rust-colored or brownish yellow, and others in which the soil is medium sand or very fine sand in texture. The soil is free from gravel, and litmus-paper tests indicate that the surface soil is acid. In its virgin condition the surface inch or two usually contains sufficient organic matter to give the soil a rather dark color, but when cultivated this becomes oxidized and the lighter color prevails.

Areas of Plainfield fine sand occur in all the townships in the western half of the county, and also in Berlin and Manchester Town-

ships. There are no large tracts of this soil but many small irregular patches ranging in size from a few acres to half a square mile.

This land is level or gently sloping, and occurs principally on stream terraces and on small islands in marshland parts of the county. The elevation of these islands in the marshland is generally only a few feet higher than the marsh, and in such places the water table is near the surface. Because of the sand in soil and subsoil, crops suffer for lack of moisture during dry seasons. Areas adjoining marshland suffer least in this respect, but early in the spring they are sometimes too wet.

Much of this land has been cleared and placed under cultivation, but much that has been cleared is no longer used as plowland. A large acreage of this soil supports a growth of brush, scrub oak, and jack pine. The chief crops are rye, potatoes, corn, and beans, and a small acreage is in oats and hay. Considerable difficulty is experienced in obtaining successful stands of grass, because the soil dries out, it is in need of lime, and is rather poor. Yields of all crops are low, except where attention is paid to fertilization, liming, and rotation. The same methods of improvement suggested for Plainfield fine sand apply to this soil.

#### SUPERIOR FINE SANDY LOAM

Superior fine sandy loam, to a depth of 8 or 10 inches, is brown medium or fine sandy loam. The subsoil is light-brown or pinkish fine sand material underlain at depths ranging from 18 to 30 inches by the red clay substratum characteristic of the Superior soils. The depth to this heavy clay may range from 1 to 3 feet, but it is everywhere present. The color and texture of the surface soil may also vary but it is everywhere light colored and sandy, being in some places sand or fine sand.

Areas of this soil are small and occur on lowlands along Fox River, mostly in the northern part of the county near Berlin. A few patches, some of which are too small to be mapped, occur at other places close to the river in the western and southwestern parts of the county.

This land is level or nearly so, and because of its sandy nature drainage is usually good. In a few areas where clay is close to the surface, drainage is somewhat deficient; as a rule, tile drains are not needed, but in some places they would be beneficial. This soil is used principally for general farming, but it is better adapted to truck crops and intensive farming.

#### MAUMEE SILT LOAM

The topsoil of Maumee silt loam is dark-brown or black silt loam, which contains a large quantity of organic matter. This is underlain at a depth of 10 or 12 inches by dark-colored silt loam, which, with increase of depth, becomes grayish-drab or mottled silt loam material grading at a depth of about 2 feet, into a mottled yellow, drab, and brown, fine sandy substratum. The material composing the substratum continues to a depth of more than 3 feet, and in places contains thin seams of clay or silt. A common variation occurs in soil with a covering of peat, several inches thick, over the surface soil. In a few small areas this peat layer, as a result of






FIG. 1.—HEMP MILL AT MARKESAN, WIS.




FIG. 2.—AN AREA OF CLYDE SILT LOAM USED FOR PASTURE





recent inundations, has been covered with a silt loam deposit. In some places the color is brown or yellowish brown because the soil is lower in organic matter. This soil resembles Clyde silt loam in color and drainage, but differs by having sand rather than heavy material in the substratum.

Areas of Maumee silt loam occur on lowlands along Fox River in the northern and western parts of the county, the most extensive being in the townships of Berlin, Seneca, St. Marie, and Marquette. This land is level or slightly depressed, and drainage is poor. Some of it is inundated during times of high water; and on areas near Fox River the water table is within 2 or 3 feet of the surface during much of the time.

Most of this land is now treeless, being covered with coarse grasses some of which along Fox River are of the marsh varieties. The original tree growth consisted of soft maple, elm, willow, ash, alder, and some moisture-loving shrubs. There is very little merchantable timber on this land.

Because this land is low and wet, not much of it has been improved. On some areas bordering the upland and along well constructed ditches where drainage is sufficient, crops can be grown. On such areas yields are good, but most of this soil is not well drained and is suitable only for hay and pasture. This soil, especially where it is deepest, is very good and when drained is suited to corn, hay, cabbage, onions, and root crops. Proper drainage is the first and most important step in the improvement of Maumee silt loam, but no great improvement can be made at present because of the Government locks on Fox River. The construction of ditches and drains where the present fall is sufficient is the only means of improving drainage at present. The land which can not be drained can be used for hay production and pasture.

#### MAUMEE LOAM

The topsoil of Maumee loam, to a depth of 10 or 12 inches, consists of black loam, containing a large quantity of organic matter. The subsoil is mottled gray, drab, or yellowish fine sand to depths of 4 or more feet, and in places contains lenses of silt and clay. The surface soil in many places contains considerable fine sand and in a few places this material is covered with several inches of peat which becomes mixed with the mineral soil material upon cultivation and soon disappears. Underlying some of the Maumee loam, as in sections 10 and 12 in Kingston Township, is a bed of pinkish-red clay material similar to that which forms the subsoil of the Superior soils. Some patches of Maumee fine sandy loam are included in mapped areas of this soil.

Areas of Maumee loam cover 12.4 square miles. This soil occurs on lowlands along Fox River in the northern and western parts of the county in Berlin, Seneca, St. Marie, Princeton, and Marquette Townships, and along Grand River in the southern part. The land is level, and very poorly drained. In its present condition it is used chiefly for pasture, and some hay is grown. Areas bordering the river are very wet and are covered with flood waters for long periods

during spring and early summer, partly on account of the locks in Fox River which retard the drainage.

Much of the land is treeless, but in some places the growth consists of soft maple, elm, ash, alder, and willows. Practically all merchantable timber was removed long ago. The chief use of this soil is for hay production and for pasture, although small patches are under cultivation. When properly drained it is a good soil for corn, hay, and truck crops, such as cabbage and onions. Grains can be grown, but there is danger of lodging.

Any great improvement in drainage on the lower-lying areas will be difficult or impossible so long as the locks remain in the river, but some of the land lying at a distance from the river and bordering the highland has sufficient fall so that under present conditions it can be successfully drained.

#### CLYDE SILT LOAM

The topsoil of Clyde silt loam consists of a nearly black, smooth silt loam very high in organic matter. This is underlain at a depth of 10 or 12 inches by silt loam material which is somewhat lighter in color in the upper part of the layer, but gradually changes, with depth, through dark grayish-brown, to drab, bluish, or yellow mottled silty clay loam material, which continues to a depth of more than 3 feet. In places the lower part of this layer contains small lenses of fine sand. This soil is similar to Maumee silt loam, except that the Maumee soil has sandy subsoil.

The chief variation in Clyde silt loam is a covering of peat which in some places reaches a depth of 8 inches. When such land is cultivated the underlying mineral soil is mixed with the peat, so that after a few years the material has the character of a mineral soil.

Clyde silt loam occurs chiefly in the eastern half of the county in depressions in areas of Miami and Parr silt loams. The largest area is in the northeastern part of Mackford Township. Smaller areas are mapped in Green Lake, Brooklyn, and Manchester Townships.

The surface of Clyde silt loam is low, level, nearly level, or slightly depressed. The drainage is poor, and some form of artificial drainage is necessary before general farm crops can be successfully grown. Tile drains have greatly increased the producing power of this soil.

The native vegetation on this soil consisted of elm, ash, willow, soft maple, and grasses. All the timber has been removed and most of the areas are cleared and cultivated or used for pasture (Pl. L, fig. 2).

Where well drained this soil makes excellent farming land, and is considered the best corn land in the southern part of the State. It is also well suited to the production of hay and root crops. Small grains have a tendency to lodge, and the quality is not so good as that of those grown on light-colored upland soils. If corn is grown for two or three years in succession the grain crop following is not so apt to lodge as it is if it follows corn grown for only one year.

#### CLYDE LOAM

Clyde loam includes dark-colored, low, poorly drained land, where the surface texture is fine sandy loam or loam and the subsoil is

gray or mottled yellow and gray sandy clay. Lenses and strata of fine sand are present throughout the soil. The soil has the same general characteristics, similar agricultural value, and occurs under practically the same conditions as Clyde silt loam. When thoroughly drained it would be well suited for the production of corn and truck crops.

Suggestions for the improvement of this soil are similar to those for Clyde silt loam.

#### POYGAN SILT LOAM

Poygan silt loam is very dark grayish-brown silt loam to depths ranging from 10 to 16 inches, containing much organic matter, and underlain by a red clay subsoil. On some small areas, a layer of peat covers the surface. In places, the subsoil is blue clay instead of the typical red clay.

Areas of Poygan silt loam occur on lowlands along Fox River in Berlin, Seneca, and St. Marie Townships. Natural drainage is poor, and tile or other drains are necessary before the soil can be successfully cultivated. When drained it is good soil for corn, hay, and root crops.

#### POYGAN CLAY LOAM

Poygan clay loam, to a depth of 6 or 8 inches, consists of heavy nearly black clay loam, grading downward into gray clay, which, at depths varying from 12 to 24 inches, is underlain by a heavy compact, pinkish-red clay, similar to that underlying soils of the Superior series. In places, a thin layer of peat covers the surface, but after a few years of cultivation this disappears. In some places, the subsoil is a heavy blue clay instead of red clay, but the texture and structure remain the same, the difference in color being due to differences in degree of oxidation. Where drainage conditions are very poor the subsoil is mottled. Thin seams of sand occur in the subsoil in a few places.

Poygan clay loam is of small extent in Green Lake County, and is confined to the valleys of Fox River and its tributaries. It occurs in Berlin, Brooklyn, St. Marie, Kingston, and Marquette Townships and covers a total area of 4.2 square miles.

The surface of this soil is low and level and the natural drainage is very poor. It occupies some of the lowest land along Fox River and must be drained before cultivated crops can be safely grown. The original water-loving trees and grasses included soft maple, willow, elm, and ash. Most of the trees have been cut and removed.

Only a small part of this land has been cleared, drained, and cultivated. When thoroughly drained the soil is fertile and well adapted to a variety of crops, including corn, hay, sugar beets, cabbage, and root crops. Small grain is apt to lodge, and the quality of the grain is not so good as that of grain grown on light-colored upland soils. Tile drains should be installed, as practically all the soil is so situated that it can be successfully drained.

#### DUNNING FINE SAND

The surface soil of Dunning fine sand, to a depth of about 4 or 6 inches, consists of black fine sand which contains a high percentage of vegetable matter. In places there is sufficient organic matter to



make the soil peaty or mucky sand. This dark material grades into dark-gray fine sand, which in places becomes gray or nearly white and is stratified in the lower part of the subsoil. This light-colored material continues to a depth that is undetermined but always greater than 3 feet. The color of the surface soil varies from black to brown, yellowish, or mottled brown and yellow, owing in part to poor drainage and in part to iron stains. In a few places there is sufficient clay in the deep subsoil to make it somewhat sticky. The surface soil in places is 14 inches deep. Included with mapped areas of this soil are a few small areas of Dunning sand, which is a soil type gradational between Plainfield sand and shallow peat.

Dunning fine sand is widely distributed in the northern and western parts of the county. It is confined entirely to the lowlands bordering Fox and Grand Rivers. The largest area, a continuous tract of several square miles, is immediately north of Lake Puck-away in Marquette Township. Most of the other areas are much smaller and in places border areas of peat.

This land is level, low, and naturally very poorly drained. Some of it is sufficiently high so that water seldom covers the surface, but in other places from 1 to 3 feet of water may cover it part of the year. Much of this land lies so low and so close to the river that drainage will be difficult unless the Fox River locks are removed.

The original growth consisted of willows, elm, soft maple, and coarse marsh grasses, but at present much of this land is open and treeless and covered with grass, cat-tail, and rushes, but on the driest places, some oak, jack pine, poplar, and white birch are present.

Only a small part of this soil is under cultivation, and crops usually are poor because of excessive moisture in the soil. Alsike clover, wild hay, and corn are the principal crops. Some potatoes are grown on the better-drained areas.

This soil is of low agricultural value. Although the surface soil is well supplied with organic matter, it is deficient in mineral plant food elements, is too sandy for most crops, and is acid. Drainage is the first step toward its improvement, and this may be so difficult, that the expense would not be justified under present conditions.

#### GENESEE SILT LOAM

The topsoil of Genesee silt loam is dark brown, and the subsoil material is brown or yellowish brown. The texture ranges from heavy fine sandy loam to silt loam. Areas occur as narrow strips of alluvial land along streams, and are subject to annual overflow.

This land has low agricultural value, but in a few places it is used for grazing. It is doubtful if it is possible to improve this soil at present, because of the high cost of artificial drainage and protection from overflow. Where improved, it is good farming land, especially for such crops as hay and corn.

#### PEAT

The material classified as peat consists chiefly of accumulated vegetable matter in varying stages of disintegration and decomposition, having more or less mineral matter mixed with it. In color, peat varies from brown to black. The depth of the material forming this soil is also extremely variable, typical peat being more than 18



inches deep and ranging from that depth to as much as 15 feet. Shallow peat ranges from 6 to 18 inches deep. Usually the shallow peat is more thoroughly decayed, and is of darker color. The mineral material underlying most of the peat is fine sand.

The color of peat and the stage of its decomposition are also variable. These variations are important, but they are not indicated on the soil map, except as they are indicated by the depth of the peaty material. Most deep peat is brown, and the structure is fibrous, which shows that it has not reached an advanced stage of decomposition. In its rawest condition the stems, leaves, grasses, and moss from which it was formed are still visible. This raw peat, in many places, continues to depths ranging from 3 to 6 feet or more, but usually peat in its lower depths is more decayed and darker than that nearer the surface. In some places deep peat is so thoroughly decomposed that its fibrous character has disappeared. Peat of this kind has more agricultural value than raw peat. In general, shallow peat is somewhat more decayed and of darker color than deep peat, and in some places it approaches muck in composition. Areas where the peat is well decomposed are small.

The mineral matter under the peat beds consists chiefly of white or grayish fine sand; but where areas of peat adjoin areas of Bellefontaine fine sandy loam, dark-colored phase, and Bellefontaine loam, the underlying layer is sticky, stony material, or sandy clay loam material. In the southeastern part of the county, small areas of peat occur along some of the drainage ways, where surrounding upland soils are heavy. On such areas the material underlying the peat is heavy in texture.

Peat is the most extensive soil in Green Lake County, occurring to some extent in every township. The largest areas are in the northern and western parts of the county in the lowlands adjacent to Fox River. Areas of deep peat are more extensive than those of shallow peat, the largest being along White River in Seneca Township, along Fox River in Princeton Township, and along Grand River in Marquette and Kingston Townships. Areas of shallow peat occur throughout the same regions as deep peat and in many places they border large marshes. Shallow peat may be considered a gradational soil between typical peat and soils of the Dunning series.

Tracts of peat soil are comparatively low, flat, and very poorly drained. Water stands on the surface of many of the marshes during spring and early summer, and when in this soggy condition the land is so soft that it will not support the weight of cattle. During late summer, especially in dry seasons, the marshes dry out so that farm animals are safe, and the peat frequently becomes so dry that danger from fire is great. All soils included in mapped areas of peat are sufficiently high in organic matter to burn when dry.

Many large drainage ditches have been extended into and through the peat areas, but these supply only partial outlets, so that numerous lateral ditches supplemented by tile drains are necessary before crops can be profitably grown. The drainage of most peat areas along Fox River is obstructed by dams in the river.

The forest growth on peat marshes consists of tamarack, alder, poplar, willow, and various other water-loving trees. Only a small acreage of peat marsh is forested, most of it being open and treeless.

On open marshes there is a growth of coarse marsh grass, wire grass, or sphagnum moss, interspersed with a small and stunted growth of water-loving shrubs. Some of the marsh grasses are pastured or cut for hay.

Although areas of peat are extensive in Green Lake County, this land is at present of little agricultural importance. Some cultivation is being attempted, but in most places drainage is inadequate, and crop failures have resulted because of an excess of moisture on land which produced fair or good crops of timothy hay during a series of dry seasons. Commercial fertilizers and lime are needed, but are not used very much on marshes now under cultivation.

Crops most commonly grown on peat are buckwheat, rye, timothy, potatoes, and root crops; and some cabbage and onions. Some attempts are made to grow corn, but because of the danger of summer frosts this crop is uncertain.

Some tracts of peat are utilized to a small extent for pasture and hay, although wild marsh grasses have low feeding value. These marshes are frequently burned over to destroy the dead grass and trash upon the surface, and a moderate stand of clean grass usually follows, which makes fairly good pasture while it is young and tender. If the marshes are burned over during dry seasons the peat itself is in danger of being burned.

Yields of crops mentioned, when grown without fertilization on raw, brown, fibrous peat, are usually low and unsatisfactory, but where peat is well decomposed and black fair crops may be obtained for a few years without fertilization. On some areas where a few inches at the surface have been burned sufficient concentration of mineral elements prevails to insure two or three fair crops. The fire, if uncontrolled, however, may be more detrimental than beneficial, for deep holes may be formed and the surface of the ground lowered to such an extent that the land will no longer be sufficiently drained.

It is well known that frosts frequently occur on marshland when there is none on higher land. This is partly because the cold air which forms on the surface of all ground at night, tends to flow down and collect in low places; but it is also because the loose, spongy soil of peat marshes does not conduct the sun's heat downward, and consequently, the material beneath the surface does not become warmed as in mineral soil. The little heat absorbed by the surface is rapidly lost at night by radiation, so that the freezing point is frequently reached on peat when it would not be on mineral soils.

This difficulty with peat marshes can be overcome somewhat by heavy rolling, which, by compacting the soil, enables it to conduct heat downward more readily. It will also tend to hasten decomposition, causing the peat to take on more quickly the character of muck. Killing frosts may occur on marshland at any given point as early as on upland soil having good air drainage, situated 150 miles farther north. Frost may occur two weeks or more earlier on the marshland regions of Green Lake County than on the hilltops. This means that corn and potatoes, although safe crops for the upland region, are not safe crops for the marshland.

Peat soils are comparatively low in phosphorus, potassium, calcium, and magnesium, but high in nitrogen and organic matter. The

average phosphorus content of peat in this region, according to analyses, is 0.135 per cent. This means that to a depth of 1 foot, an acre of soil contains approximately 675 pounds of phosphorus.

The deficiency of potassium in this soil is greater than that of phosphorus. It contains an average of 0.3 per cent of potassium, whereas good upland clay loam soils average 2 per cent. The average potassium content in the upper 2 feet of upland soils is approximately 120,000 pounds per acre, whereas peat soils contain only about 3,000 pounds.

The large quantity of organic matter in peat soil increases the supply of nitrogen. This soil averages 2.5 per cent of nitrogen, whereas the upland silt loam soils contain about 0.12 per cent in the 8 inches nearest the surface.

The rational treatment of peat lands requires the use of fertilizers containing phosphorus and potassium. These elements are contained in comparatively small quantities in barnyard manure, and heavy applications of manure will secure good yields of crops on peat, but manure contains much nitrogen which is not needed on the peat. Therefore, when a farm includes upland soils as well as peat, the manure should be used on the upland soils, and commercial fertilizers containing phosphorus and potassium should be used on the peat land.

On deeper peat soil which is in a raw and acid condition, the use of lime in addition to commercial fertilizers will be profitable. On a few marshes the nitrogen in the organic matter does not become available to crops, because of the low soil temperature and the high degree of soil acidity. In such cases a light application of composted stable manure, to inoculate the soil with the proper organisms, is very helpful.

Crops recommended for peat are grasses for hay and pasture, hardy root crops, rye, and oats. When the land is properly fertilized and limed, clover, alfalfa, and other legumes can be grown. On fairly well drained, decomposed marshland good pasture can be developed, as the compacting of the soil by livestock is very beneficial; and where the pasture is good and other conditions are favorable, certain parts of these marshes can be successfully utilized for dairying or stock raising. When peat land is cultivated, a heavy roller should be used to compact it. Certain special crops, such as cabbage, onions, buckwheat, and rape, are adapted to properly fertilized and well-drained peat land.<sup>3</sup>

In summarizing the prospects for the future agricultural development of peat land in this county, several conditions obtain with which it is necessary to comply before farming on this land can be permanently successful.

1. It is absolutely necessary that the land be sufficiently drained. Large outlet ditches, although necessary, are not in themselves sufficient, but must be supplemented by open laterals and tile drains.
2. Since this land is generally deficient in potash, phosphorus, and lime, these materials must be supplied in proper form and adequate quantities to insure permanent profitable production.
3. It must be recognized that danger from summer frosts makes such crops as corn and potatoes uncertain, and the crops to be grown

<sup>3</sup>For a more complete discussion of the management of marsh soils see the following bulletin: WHITSON, A. R., and SIEVERS, F. J. THE DEVELOPMENT OF MARSH SOILS. Wis. Agr. Expt. Sta. Bul. 205, 22 p., illus. 1911.



must be suited not only to the soil, but also to the climatic conditions.

4. The use of a heavy roller, to compact the soil, is necessary in the cultivation of peat land.

5. Those purchasing peat land for farming purposes must see their way clear not only to pay for the land but also to provide adequate drainage and fertilization, both of which call for further outlay of money.

It is suggested to those who desire to undertake the development of a peat farm but who have had no experience with this kind of land, that they rent such a farm for a year or two, or serve an apprenticeship on a farm where peat is successfully farmed, thus acquiring valuable experience at minimum cost.

*Peat, shallow phase.*—Areas of peat where the organic material is less than 18 inches thick are indicated separately on the soil map as peat, shallow phase. This phase of peat is described in the foregoing general discussion of peat soils.

#### ROUGH STONY LAND

Rough stony land in Green Lake County includes land which is so stony, rocky, steep, and broken that it can not be cultivated. Some of this land can be used for grazing, but much of it consists of rock outcrop on which but small patches of soil occur. Some of the areas are steep cliffs of outcropping rocks, and other tracts are entirely stones or rock. Some of the land is not very rough, but the rock is so near the surface as to render the land of little agricultural value. On a few areas the rock has a fair covering of soil, but the land is so steep that it is impossible to cultivate it, though it is used for grazing. Soil which occurs on rough stony land ranges from heavy to light in texture, depending upon the nature of the rock.

Limestone, sandstone, and granite are present in rough stony land of this county. The eastern part of the county is generally underlain by limestone and the western part by sandstone. Projecting through these formations are granite gneiss and other crystalline outcrops in the vicinity of Utley, within the city limits of Berlin, and immediately south of Lake Puckaway.

The only agricultural value of rough stony land is in the little pasturage which it affords. In a few places cultivation is attempted, but it is carried on with difficulty and with unsatisfactory results. Where steep slopes are forested they should be allowed to remain so, and where timber has been removed, reforestation should be considered.

At Utley and Berlin, rock is quarried from the granitic outcrops. At Berlin a very good quality of rock is present, some of which is used for monuments. Building stone, paving stone, curbstone, and some road material are among the products of the quarries.

#### SUGGESTIONS FOR SOIL IMPROVEMENT

##### LIME AND FERTILIZERS

*Liming.*—On the prairies the topsoils are acid and the subsoils show some acidity to depths varying from 2 to 3 feet. The light-colored heavy upland soils are usually slightly acid at the surface,



but the material of the gravelly layer in the subsoil contains carbonate of lime. The sandy soils are most in need of lime, and the soils least in need of lime are Clyde, Poygan, and Superior soils.

The degree of acidity is variable and each farmer may find a wide variation in soil acidity on his farm. Failure of clover and alfalfa to make a stand is a good indication of the need of lime. Such crops as alfalfa, sweet clover, peas, cabbage, onions, and lettuce have a high lime requirement; clover, garden beans, barley, hemp, turnips, and radishes have a medium lime requirement; and vetch, white clover, oats, rye, bluegrass, potatoes, sorghum, and some others have a low lime requirement. As a rule the heavy soils which are acid need more lime than the sandy ones which show the same degree of acidity.

Ground limestone is doubtless the most economical form of lime for Green Lake County. Lime should be applied before planting the crop which is to be benefited. It should be applied to plowed land and thoroughly worked in by harrowing. Either fall, winter, or spring applications may be made on heavy soils, but on light soils spring applications are preferable. The best way to apply lime is with a lime spreader, but a manure spreader may be used by putting a thin layer of manure over the bottom of the spreader and then spreading the limestone evenly on top of it. Where several farmers are so situated that they can work together a lime spreader may be purchased jointly for this purpose.

After making an initial application of 2 tons of ground limestone per acre it is not likely that another application will be needed for about four or six years. Subsequent applications should be determined by the needs of the crops and by acidity tests.

Most acid soils are deficient in available phosphorus, and the need of phosphorus may be so great that lime has little effect until phosphorus is added. In places the application of phosphate alone to an acid soil will result in larger increases in yield than the use of lime alone, and for this reason it is important that both deficiencies should be corrected to obtain the best results.

Ground limestone may be applied at rates ranging from 2,000 to 4,000 or more pounds an acre, whereas 75 to 400 pounds of phosphates and other fertilizers are usually applied for staple crops. A machine for distributing ground limestone should be provided with a double agitator so as to effect continuous feeding. End-gate spreaders which will distribute proper quantities of either fertilizer or ground limestone fairly well are available. When a fertilizer distributor is not available, acid phosphate or other fertilizer may be spread evenly over the load of manure in the manure spreader and distributed fairly well in this manner. The quantity of fertilizer to be spread over each spreader load must be calculated so that the right quantity will be applied to the soil. An old drill or seeder may also be used to distribute fertilizer, the grain drill following the fertilizer drill.

*Use of farm manure.*—The care and use of the manure produced is an important factor in the management of livestock farms. The principal advantage of farming with livestock is that the manure produced may be used to maintain profitable yields, so that comparatively little fertilizer need be purchased. Much of the crop nutrients in manure is readily soluble in water, so that if the manure is ex-

posed to much rain a considerable part of its soluble material is lost. It is important also to remember that a large part of the nutrient elements is in the liquid part of the manure, making it necessary to use bedding or other absorbents freely to prevent a considerable loss of the fertilizing value of the manure. This is particularly true of potassium, about 60 per cent of which is contained in the liquid part.

Ordinarily the best practice is to haul the manure directly to the field. When manure must be kept in a pile for a time, the pile should be kept compact and moist. In this climate the use of shelter for manure is of doubtful importance, though where more rains occur, particularly in the winter, a shed roof is very helpful to conserve manure.

The rate and frequency with which manure is applied depends in part on the character of the soil. On lighter soils, frequent applications of small quantities prove most effective. An application of 5 or 6 loads an acre once in three years is desirable on the sandy loams, whereas from 8 to 12 or more loads once in four or five years may be used to advantage on the heavier soils.

#### CROP ROTATIONS

Owing to the great variety of soils in Green Lake County, many kinds of crops may be grown to advantage. The kind of crop or the rotation to be followed on the sandy lands will not be suited to the black prairie lands or to the clay lands, and the system of farming followed on the marshlands will not give best results on the uplands.<sup>4</sup>

It is generally recognized that it is advisable to follow a definite crop rotation, or at least to diversify crops, rather than to have the same crop growing on a certain field year after year. It seems to be easier to maintain fertility through diversification of crops than by practicing a one-crop system of farming.

Crop rotation helps to keep down weeds, aids in controlling plant diseases, serves to check insect pests, and helps to maintain soil fertility. A good system of cropping increases the humus supply in the soil, and also aids in the distribution of farm labor, rendering it more effective.

Among the factors which determine the kind of rotation to practice are (1) the type of soil, (2) the kind and acreage of crops a farmer wishes to grow, (3) the surface features of the land, and (4) soil fertility.

The following rotation is probably the best for most sandy soils: First year, clover with perhaps a light seeding of rye; second year, clover for hay, leaving the second crop to be turned under either in fall or spring; third year, corn or potatoes; fourth year, soy beans, which may be used for seed, for hay, and for green manure. If any other crops are to be grown, they may be planted after clover.

The following rotations are suggested for heavy soils: Five-year rotations—corn, corn, small grain (with clover), clover, and wheat (with clover); or corn, small grain (with clover), clover, wheat (with clover), and clover. Four-year rotations—wheat, corn, oats

<sup>4</sup>For a full discussion of crop rotations consult the bulletins of the University of Wisconsin Agricultural Experiment Station.

(with clover), and clover; or corn, corn, oats (with clover), and clover. Three-year rotation—corn, oats (with clover), and clover.

It seems unnecessary to add organic matter to peat soil which in itself is practically all organic matter, yet it is recognized as a good practice, especially where potatoes, cabbage, sugar beets, and onions are grown. The use of proper fertilizers, combined with a crop rotation containing clover, or clover and timothy, is advisable. On farms consisting entirely of marshland it is as important to grow alfalfa and clover as it is on upland soils. With the use of a legume or hay crop in the rotation some small grain such as rye or oats must necessarily be grown, but they are considered less desirable crops for this kind of soil because of their tendency to lodge. If corn or some other cultivated crop is raised the first three years and the soil is treated with the proper mineral fertilizers during that time, small grain may be grown very successfully, provided less seed be sown than on upland soil. If the grain crop is too rank it would be well to cut it for hay in order to have the grass or clover seeding.

If a farmer has 30 acres of peat and desires to grow cabbage and sugar beets as cash crops, it is a good plan to divide the field into four strips and crop them to sugar beets, cabbage, grain, and hay, so that a four-year rotation of hay, sugar beets, cabbage, and grain would be practiced on each strip. On a dairy farm, two or three corn crops may be grown in succession, to be followed by small grain, and then seeded to alsike clover and timothy. The hay may be cut the first year and pastured the second. The rotation best suited to any peat marsh can best be determined by the conditions existing on that particular farm.

A three-year rotation is popular on many of the dairy farms. It consists of grain seeded to clover, and the clover followed by corn or some other cultivated crop. Much the larger part of the corn in this region is put into the silo to be fed to dairy animals, and the clover is cured for hay and fed to livestock. It is often possible to get two crops of medium red clover in one season, the second of which may be cut for hay, or may be pastured. This three-year rotation may be lengthened into a four-year rotation by the addition of timothy, so that hay may be cut two years instead of one year, or the land may be pastured the second year instead of cutting the timothy for hay.

Potato raising, when properly managed, is profitable. Although good crops may be grown on heavy soils, the sandy loams are especially well adapted to potato production. For best results, this crop should be grown in rotation with other crops, and should always follow a legume of some kind. Potatoes should not follow corn, and corn should not follow potatoes, as both crops draw heavily on the plant-food elements of the soil. In the rotations described, potatoes may be planted as one of the cultivated crops. It is better to apply manure to the clover crop rather than to potatoes, because potato scab is more common where potatoes are planted on freshly manured land. The three-year rotation just described is excellent for sections where potatoes are grown extensively, the potatoes taking the place of corn. As a rule, cropping to potatoes more frequently than once in three years is not recommended.



As indicated elsewhere, hemp growing is important in this county. Regarding the best crops to grow in rotation with hemp, the following may be said:<sup>5</sup> Hemp should not follow timothy meadow, bluegrass sod, nor pasture. Neither should hemp follow any small-grain crop, unless the soil is very well supplied with manure.

Hemp leaves the soil in excellent physical condition for any spring-sown small grain. It also leaves the land comparatively free from weeds, and consequently it is a good crop to precede sugar beets or canning peas, for these crops especially require a clean seed bed. The following rotations have given good results in growing hemp:

Four-year rotation—small-grain crop (seeded with clover); clover for hay and pasture (manured and fall plowed); corn, potatoes, or similar crop; and hemp (then back to small grain and clover). In this rotation many growers sow timothy with the clover. There is no objection to this if the land is not left in sod for more than a year.

Three-year rotation—small-grain crop (seeded to clover); clover (manured and fall plowed); hemp (then back to small grain and clover). In this rotation considerable attention should be given to using well-pulverized manure and to spreading it evenly. The clover sod should be worked thoroughly in the spring to insure a compact, smooth seed bed.

#### SUMMARY

Green Lake County is slightly southeast of the geographical center of Wisconsin. It has an area of 360 square miles, or 230,400 acres. The county is almost entirely within the Fox River drainage basin.

Most of the land is gently rolling, comprising level river terraces, flat marshland, gently rolling prairies, and some rolling and gently rolling clay lands and sandy loams. There is very little land too steep for cultivation.

The first settlement was made between 1830 and 1835, and the county was organized in 1858 from territory which was previously a part of Marquette County. The population in 1920 was 14,875. Green Lake is the county seat, and had a population of 456 in 1920. Berlin, with a population of 4,356, is the largest city.

Most of the county is well supplied with railroad facilities, but a few places are 9 miles by wagon road from a shipping station.

The eastern part of the county contains some of the finest prairie land in the State, but the western part contains large areas of sandy and marshy lands. The agricultural development on these different soils is markedly different, and there is a wide range in soil conditions and agricultural practices within the limits of this small county.

The soil-forming materials of Green Lake County originated from calcareous glacial drift, sandy glacial drift, water-laid materials, and from accumulated vegetable matter in varying stages of decomposition. The 31 soil types mapped are grouped into 14 series, not including peat and rough stony land.

The various soils are described in detail and their agricultural values are discussed. Some suggestions are offered for the improvement of these soils.

<sup>5</sup> See footnote 2, p. 1811.



# Accessibility Statement

---

This document is not accessible by screen-reader software. The U.S. Department of Agriculture is committed to making its electronic and information technologies accessible to individuals with disabilities by meeting or exceeding the requirements of Section 508 of the Rehabilitation Act (29 U.S.C. 794d), as amended in 1998. Section 508 is a federal law that requires agencies to provide individuals with disabilities equal access to electronic information and data comparable to those who do not have disabilities, unless an undue burden would be imposed on the agency. The Section 508 standards are the technical requirements and criteria that are used to measure conformance within this law. More information on Section 508 and the technical standards can be found at [www.section508.gov](http://www.section508.gov).

If you require assistance or wish to report an issue related to the accessibility of any content on this website, please email [Section508@oc.usda.gov](mailto:Section508@oc.usda.gov). If applicable, please include the web address or URL and the specific problems you have encountered. You may also contact a representative from the [USDA Section 508 Coordination Team](#).

## **Nondiscrimination Statement**

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the

Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.




To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [http://www.ascr.usda.gov/complaint\\_filing\\_cust.html](http://www.ascr.usda.gov/complaint_filing_cust.html) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by:

- (1) mail: U.S. Department of Agriculture  
Office of the Assistant Secretary for Civil Rights  
1400 Independence Avenue, SW  
Washington, D.C. 20250-9410;
- (2) fax: (202) 690-7442; or
- (3) email: [program.intake@usda.gov](mailto:program.intake@usda.gov).

USDA is an equal opportunity provider, employer, and lender.



CONVENTIONAL  
SIGNS



LEGEND

Clyde loam	Miami loam
Cy	Mi
Clyde silt loam	Miami silt loam
Cs	Ms
Maumee loam (Clyde)	Shallow phase
Ma	Me
Maumee silt loam (Clyde)	Bellefontaine gravelly loam (Miami)
Me	Bm
Peat	Bellefontaine gravelly loam (Miami)
P	Bg
Shallow phase	Bellefontaine sandy loam (Miami)
Po	Bf
Poynan silt loam	Dark colored phase
Po	Bf
Poynan clay loam	Bellefontaine fine sandy loam (Miami)
Pc	B
Fox sandy loam	Bellefontaine silt loam (Miami)
Fs	Bs
Fox loam	Parr fine sandy loam (Carrington)
Fl	Pr
Fox silt loam	Parr loam (Carrington)
Fm	Pi
Superior fine sandy loam	Gravelly phase
Sf	Pm
Coloma sand	Parr silt loam (Carrington)
Cd	Pm
Coloma fine sand	Level phase
Cf	Wf
Dunning fine sand	Warsaw fine sandy loam (Waukesha)
Df	Ws
Dodgeville silt loam	Warsaw loam (Waukesha)
Ds	Ws
Plainfield sand	Warsaw silt loam (Waukesha)
Ps	Ws
Plainfield fine sand	Genesee silt loam
Pf	Gs
Rough stony land	
R	

